Indexes

Author Index

Abbas, R. see Kidron, M.

Abbas, Z.G. see Morbach, S.

Abian, J. see Flores, L.

Abrahamian, H. see Kästenbauer, T.

Adams, D. see Gill-Randall, R.J.

Adegoke, O.J. see Oli, J.M.

Adriaanse, M.C., Dekker, J.M., Spijkerman, A.M.W.,

Twisk, J.W.R., Nijpels, G., van der Ploeg, H.M., Heine, R.J.

& Snoek, F.J. Health-related quality of life in the first year

following diagnosis of Type 2 diabetes: newly diagnosed patients in general practice compared with screening-detected patients.

The Hoorn Screening Study, 1075

Adriaanse, M.C., Snoek, F.J., Dekker, J.M., Spijkerman, A.M.W., Nijpels, G., Twisk, J.W.R., van der Ploeg, H.M. & Heine, R.J. No substantial psychological impact of the diagnosis of Type 2 diabetes following targeted population screening: The Hoorn Screening Study, 992

Agrawal, A. see Bennett, S.M.A.

Ahmad, Y. see O'Hare, J.P.

Ahn, C.W. see Kim, C.S.

Ahn, C.W., Song, Y.D., Nam, J.H., Kim, D.M., Woo, S.O.,

Park, S.W., Cha, B.S., Lim, S.K., Kim, K.R., Lee, J.H., Lee, H.C. & Huh, K.B. Insulin sensitivity in physically fit and unfit children of parents with Type 2 diabetes, 59

Aiba, N. see Watanabe, A.

Aiso, Y. see Nagasaka, S.

Aitken, J.M. see George, A.M.

Akhter, J. see Jabbar, A.

Alberti, K.G.M.M. see Zimmet, P.

Alcolado, J.C. see Gill-Randall, R.J.

Allagoa, B. see Harvey, J.N.

Allain, T.J. see Gregson, C.L.

Allen, T.J. see Jerums, G.

Al-Mrayat, M., Samarasinghe, Y., Treml, H., Munro, N., Shotliff, K., Mcintosh, C. & Feher, M.D. A new cause of neuroglycopenia: 'missing the point', 497

Amano, S. see Mimura, T.

Amiel, S.A. see Evans, M.L.

Amit, M. see Rachmani, R.

Andaluz, P. see Audí, L.

Andersen, H. see Leinonen, A.

Andersen, H. see Nielsen, J.F.

Ando, H. see Tsuchiyama, N.

Antoniades, C., Tousoulis, D., Tountas, C., Tentolouris, C., Toutouza, M., Vasiliadou, C., Tsioufis, C., Toutouzas, P. & Stefanadis, C. Vascular endothelium and inflammatory process, in patients with combined Type 2 diabetes mellitus and coronary atherosclerosis: the effects of vitamin C, 552

Anwar, A. see Valsamakis, G.

Apfalter, P. see Daxboeck, F.

Araie, M. see Mimura, T.

Arakawa, Y. see Watanabe, A.

Arbit, F. see Kidron, M.

Armstrong, D. see Menon, V.

Arnqvist, H. see Schölin, A.

Aron.on, D., Bartha, P., Zinder, O., Kerner, A., Shitman, E., Markiewicz, W., Brook, G.J. & Levy, Y. Association between fasting glucose and C-reactive protein in middle-aged subjects,

Assaloni, R. see Ceriello, A.

Assaloni, R. see Da Ros, R.

Athwal, S. see Wood, D.M.

Atkinson, M. see Gale, E.A.M.

Attali, C. see Varroud-Vial, M.

Attali, J. see Varroud-Vial, M.

Attali, J-R. see Cosson, E.

Audí, L., Martí, G., Esteban, C., Oyarzabal, M., Chueca, M., Gussinyé, M., Yeste, D., Fernández-Cancio, M., Andaluz, P. & Carrascosa, A. VDR gene polymorphism at exon 2 start codon (Fokl) may have influenced Type 1 diabetes mellitus susceptibility in two Spanish populations, 393

Avery, A.J. see Naushahi, M.J.

Baba, T. see Fujiwara, M.

Baba, T., Østerby, R., Neugebauer-Baba, S., Nozawa, Y., Watanabe, T., Sakurai, K., Katoh, T. & Watanabe, T. No nephropathy in Type 2 diabetic patient with POEMS syndrome with an elevated plasma VEGF, 292

Bachinger, A. see Forst, T.

Bagust, A. see McEwan, P.

Bagust, A. see Shearer, A.

Baik, S.H. see Kim, N.H.

Baillet-Blanco, L. see Rigalleau, V.

Bailleul, F. see Nobels, F.

Bain, S.C. see Tait, K.F.

Baines, P.S. see Ellis, J.D.

Baker, J.R. see Dissanayake, A.M.

Baker, R. see Wilson, A.

Balkau, B. see Heine, R.J.

van Ballegooie, E. see Dijkstra, R.F.

Banerjee, A.K. see Valsamakis, G.

Banerji, M.A. see Reboussin, D.M.

Baraldi, L. see Marchesini, G.

Barbe, M. see Rigalleau, V.

Bargero, G. see Bruno, G.

Barini, A. see Spallone, V.

Barnett, A. see Valsamakis, G.

Barnett, A.H. see O'Hare, J.P.

Barnett, A.H. see Tait, K.F.

Barnett, A.H. Reply, 1058

Bar-On, H. see Kidron, M.

Barrett-Connor, E. see Palinkas, L.A.

Barry, M.J. see Grant, R.W.

Bartha, P. see Aronson, D.

Baschetti, R. Preventing Type 2 diabetes: an evolutionary view, 649 Bates, C.J., Lean, M.E.J., Mansoor, M.A. & Prentice, A. Nutrient

intakes; biochemical and risk indices associated with Type 2 diabetes and glycosylated haemoglobin, in the British National Diet and Nutrition Survey of people aged 65 years and over, 677

Bates, P.C. see Schernthaner, G.

Beauvieux, M.C. see Rigalleau, V.

Beckers, F. see Nobels, F.

Bek, T. see Luzio, S.

Ben Yehuda, A. see Eldor, R.

Benbow, S.J. see Daousi, C.

Ben-Haroush, A., Yogev, Y. & Hod, M. Epidemiology of gestational diabetes mellitus and its association with Type 2 diabetes, 103

Bennett, S.M.A., Agrawal, A., Elasha, H., Heise, M., Jones, N.P., Walker, M. & Wilding, J.P.H. Rosiglitazone improves insulin sensitivity, glucose tolerance and ambulatory blood pressure in subjects with impaired glucose tolerance, 415

Bera, L. see Varroud-Vial, M.

Berard, L. see Nyomba, B.L.G.

Berg, G. see Pell, J.P.

Bergeron, J. see Piché, M.E.

Berghold, A. see Siebenhofer, A.

Bergis, K. see Mussell, M.

Berla, M. see Rachmani, R.

Berman, J. see Tait, K.F.

Berne, C. see Schölin, A.

Betteridge, D.J. see Thomason, M.J.

Bianchi, A. see Castelli, R.

Bingley, P.J. see Swerdlow, A.J.

Birkett, M.A. see Schernthaner, G.

Bíró, L. see Jermendy, G.

Bjerregaard, P. see Jørgensen, M.E.

Björk, E. see Schölin, A.

Blaak, E.E. see van Dam, R.M.

Blohmé, G. see Schölin, A.

Blow, M. see Miller, D.R.

Bodmer, C.W. see George, A.M.

Boel, E. see Lajer, M.

Boemi, M., Sirolla, C., Testa, R., Cenerelli, S., Fumelli, P. & James, R.W. Smoking is associated with reduced serum levels of the antioxidant enzyme, paraoxonase, in Type 2 diabetic patients, 423

Bogardus, C. see Stefan, N.

Boldur, I. see Slater, R.A.

Bolinder, J. see Schölin, A.

Bolli, G.B. see Porcellati, F.

Bonadonna, R.C. see Bonora, E.

Bonora, E., Targher, G., Formentini, G., Calcaterra, F., Lombardi, S., Marini, F., Zenari, L., Saggiani, F., Poli, M., Perbellini, S., Raffaelli, A., Gemma, L., Santi, L., Bonadonna, R.C. & Muggeo, M. The Metabolic Syndrome is an independent predictor of cardiovascular disease in Type 2 diabetic subjects. Prospective data from the Verona Diabetes Complications Study, 52

Bonovas, S., Peponis, V. & Filioussi, K. Diabetes mellitus as a risk factor for primary open-angle glaucoma: a meta-analysis, 609

Borch-Johnsen, K. see Glümer, C.

Borch-Johnsen, K. see Jørgensen, M.E.

Borch-Johnsen, K. see Lauenborg, J.

Bornovali, S. see Glueck, C.J.

Botha, J.L. see Swerdlow, A.J.

Botten, R. see Russo, A.

Boucher, B.J. & Hitman, G.A. The vitamin D receptor gene variant and physical activity predicts fasting blood glucose levels in healthy young men, 803

Bouhanick, B. see Sallé, A.

Boulton, A.J.M. see Eldor, R.

Boulton, A.J.M. see Oyibo, S.O.

Boyce, J. see Usher, D.

Braspenning, J.C.C. see Dijkstra, R.F.

Brath, H. see Kästenbauer, T.

Braun, A., Muller, U.A., Muller, R., Leppert, K. & Schiel, R.
Structured treatment and teaching of patients with Type 2 diabetes
mellitus and impaired cognitive function—the DICOF trial, 999

Breislin, K. see Oyibo, S.O. Brennan, G. see James, P.

Brook, G.I. see Aronson, D.

Brooks, A.P. see Teh, M.M.

Brown, A. see Stride, A.

Brown, A., Steel, J.M., Duncan, C., Duncan, A. & McBain, A.M. An assessment of the adequacy of suspension of insulin in pen injectors, 604

Brown, J.B. see Nichols, G.A.

Brown, P.M. see Song, S.H.

Browne, R. see Menon, V.

Bruce, N.J. see New, J.P.

Bruno, G. see Monge, L.

Bruno, G., Giunti, S., Bargero, G., Ferrero, S., Pagano, G. & Perin, P.C. Sex-differences in prevalence of electrocardiographic left ventricular hypertrophy in Type 2 diabetes: The Casale Monferrato Study, 823 Buchs, A. see Slater, R.A.

Bulsara, M.K., Holman, C.D.J., Davis, E.A. & Jones, T.W. Evaluating risk factors associated with severe hypoglycaemia in epidemiology studies—what method should we use?, 914

Bundred, P.E. see Daousi, C.

Burden, A.C. see Swerdlow, A.J.

Burden, A.C.F. Re: post-prandial hypoglycaemia due to insulin lispro or insulin aspart, 1373

Burns, J.A. see New, J.P.

Burrell, L.M. see Srivastava, P.M.

Byrne, C.D. see Holt, R.I.G.

Cagliero, E. see Grant, R.W.

Caimi, G., Sinagra, D., Scarpitta, A.M., Catania, A. & Lo Presti, R. Haematocrit in metabolic syndrome, 299

Calafiore, P. see Srivastava, P.M.

Calcaterra, F. see Bonora, E.

Cameron, J. see Meyer, C.

Campbell, D.J. see Jerums, G.

Campistol, J.M. see Flores, L.

Canani, L.H. see Costa, L.A.

Cansfield, J. see Gibney, I.

Cappellini, M.D. see Castelli, R.

Cappleman, J.M. see Raskin, P.

Cargo, M.D. see Daniel, M.

Caro, I. see Caro, J.J.

Caro, J.J., Getsios, D., Caro, I., Klittich, W.S. & O'brien, J.A. Economic evaluation of therapeutic interventions to prevent Type 2 diabetes in Canada, 1229



Carr, L. see Scuffham, P.A.

Carrascosa, A. see Audí, L.

Carrozzi, G. see Ciardullo, A.V.

Carr-Smith, J. see Tait, K.F.

Casabonne, D. see Green, J.

Casals, E. see Rodríguez-Villar, C.

Cases, A. see Flores, L.

Casillas, J.M. see Vergès, B.

Casparie, A.F. see Dijkstra, R.F.

Castelli, R., Tempesta, A., Bianchi, A., Porro, T., Ivaldi, G. & Cappellini, M.D. Unreliable estimation of HbA_{1c} due to the presence of Camperdown haemoglobin [β 104 (G6) Arg \rightarrow Ser], 377

Castleden, H.A.J. see Stride, A.

Catania, A. see Caimi, G.

Cattan, S. see Cosson, E.

Cavan, D.A. see Cheyne, E.H.

Cavarape, A. see Ceriello, A.

Cenerelli, S. see Boemi, M.

Ceriello, A. see Da Ros, R.

Ceriello, A. see Heine, R.I.

Ceriello, A., Cavarape, A., Martinelli, L., Da Ros, R., Marra, G., Quagliaro, L., Piconi, L., Assaloni, R. & Motz, E. The post-prandial state in Type 2 diabetes and endothelial dysfunction: effects of insulin aspart, 171

Cerrelli, F. see Marchesini, G.

Cester, N. see Mazzanti, L.

Cha, B.S. see Ahn, C.W.

Cha, B.S. see Kim, C.S.

Cha, D.R. see Kim, N.H.

Chan, J.C.N. see Chan, W.B.

Chan, W.B., Tong, P.C.Y., Chow, C.C., So, W.Y., Ng, M.C.Y., Ma, R.C.W., Osaki, R., Cockram, C.S. & Chan, J.C.N. The associations of body mass index, C-peptide and metabolic status in Chinese Type 2 diabetic patients, 349

Chantelau, E. A review of basal insulins, 1057

Chantelau, E. Reply, 647

Chapman, I.M. see Russo, A.

Charpentier, G. see Varroud-Vial, M.

Chaturvedi, N. see Jafar, T.H.

Chen, Y-T. see Yu, H-I.

Chetty, R. see Valsamakis, G.

Chetwynd, A. see Marshall, A.L.

Cheyne, E.H., Sherwin, R.S., Lunt, M.J., Cavan, D.A., Thomas, P.W. & Kerr, D. Influence of alcohol on cognitive performance during mild hypoglycaemia; implications for Type 1 diabetes, 230

Chien, J. see Rave, K.

Choi, D.S. see Kim, N.H.

Choi, K.M. see Kim, N.H.

Chow, C.C. see Chan, W.B.

Chueca, M. see Audí, L.

Chueh, H.C. see Grant, R.W.

Ciardullo, A.V., Fattori, G., Carrozzi, G. & Daghio, M.M. Evaluation of an information tool for diabetic patient education, 1056

Clària, J. see Flores, L.

Classen, J.B. Pertussis infections, vaccines and Type 1 diabetes, 397

Clausen, J. see Tan, M.H.

Cleland, S.J. see McEleavy, O.D.

Cockram, C.S. see Chan, W.B.

Cockram, C.S. see Pan, C.Y.

Cohen, M. see Vergès, B.

Coisne, D. see Hadjadj, S.

Coker, R. see Shapo, L.

Colagiuri, S. see Heller, S.R.

Cole, A. see Ellis, J.D.

Cole, D.R. see Simmons, D.

Coleman, M.A.G. see Holt, R.I.G.

Colhoun, H.M. see Thomason, M.J.

Coll, A.P. & Thomas, S. Diabetic ketoacidosis presenting in a patient with Addison's disease, 1155

Collins, J.E. see Tait, K.F.

Colquhoun, I. see Pell, J.P.

Compston, A. see Tait, K.F.

Conget, I. see Flores, L.

Connell, J.M.C. see McEleavy, O.D.

Connelly, P.W. see Retnakaran, R.

Cook-Mozaffari, P. see Goldacre, M.J.

Cooper, J. see Foo, K.

Cooper, M.E. see Jerums, G.

Cooper, M.E. see Mogensen, C.E.

Cosson, E., Paycha, F., Paries, J., Cattan, S., Ramadan, A., Meddah, D., Attali, J-R. & Valensi, P. Detecting silent coronary stenoses and stratifying cardiac risk in patients with diabetes: ECG stress test or exercise myocardial scintigraphy?, 342

Costa, L.A., Canani, L.H., Lisbôa, H.R.K., Tres, G.S. & Gross, J.L. Aggregation of features of the metabolic syndrome is associated with increased prevalence of chronic complications in Type 2 diabetes, 252

Coster, G. see Simmons, D.

Creagh, F.M. see Moustafa, M.

Crombie, I. see Ellis, J.D.

Crook, M. Type 2 diabetes mellitus: a disease of the innate immune system? An update, 203

Crouse III, R.J. see Reboussin, D.M.

Cull, C.A., Neil, H.A.W. & Holman, R.R. Changing aspirin use in patients with Type 2 diabetes in the UKPDS, 1368

Cummings, M.H. see Gibney, J.

Currie, C.J. see McEwan, P.

Curzi, C.M. see Mazzanti, L.

Daghio, M.M. see Ciardullo, A.V.

Dalton, R.N. see Schultz, C.J.

van Dam, R.M., Hoebee, B., Seidell, J.C., Schaap, M.M., Blaak, E.E. & Feskens, E.J.M. The insulin receptor substrate-1 Gly972Arg polymorphism is not associated with Type 2 diabetes mellitus in two population-based studies, 752

Damm, P. see Lauenborg, J.

Daneman, D. see Zdravkovic, V.

Dani, F. see Monge, L.

Daniel, M. & Cargo, M.D. Association between smoking, insulin resistance and β-cell function in a North-western First Nation, 188

Daousi, C., MacFarlane, I.A., Woodward, A., Nurmikko, T.J., Bundred, P.E. & Benbow, S.J. Chronic painful peripheral neuropathy in an urban community: a controlled comparison of people with and without diabetes, 976

Darko, D.A. see Ng, R.S.H.

Da Ros, R. see Ceriello, A.

Da Ros, R., Assaloni, R. & Ceriello, A. The preventive anti-oxidant action of thiazolidinediones: a new therapeutic prospect in diabetes and insulin resistance, 1249

Dashora, U.K. & Taylor, R. Maintaining glycaemic control during high-dose prednisolone administration for hyperemesis gravidarum in Type 1 diabetes, 298

Davies, M.J., Tringham, J.R., Troughton, J. & Khunti, K.K.

Prevention of Type 2 diabetes mellitus. A review of the evidence and its application in a UK setting, 403

Davis, E.A. see Bulsara, M.K.

Dawson, R.A. see Moustafa, M.

Daxboeck, F., Goerzer, E., Apfalter, P., Nehr, M. & Krause, R. Isolation of *Bordetella trematum* from a diabetic leg ulcer, 1247

Day, T. see Delaney, C.

Deaner, A. see Foo, K.

Dekker, J.M. see Adriaanse, M.C.

Delaney, C., Shaw, J. & Day, T. Acute, local effects of iontophoresed insulin and C-peptide on cutaneous microvascular function in Type 1 diabetes mellitus, 428

Del Prato, S. see Heine, R.I.

Democratis, J. see Saunders, S.A.

De Neeling, A.J. see Meiland, R.

De Schrijver, P. see Nobels, F.

Després, J.P. see Piché, M.E.

Devendra, D., Galloway, T.S., Horton, S.J. & Wilkin, T.J. Exploring the idiotypes of insulin antibodies as markers for remission in Type 1 diabetes, 1316

DeVries, J.H., Snoek, F.J. & Heine, R.J. Persistent poor glycaemic control in adult Type 1 diabetes. A closer look at the problem, 1263

Dewhurst, J.A. see Warburton, T.J.

Dharashivkar, S. see Glueck, C.J.

Dickey, R. see Ratner, R.E.

Dijkstra, R.F., Braspenning, J.C.C., Huijsmans, Z., Peters, S., van Ballegooie, E., ten Have, P., Casparie, A.F. & Grol, R.P.T.M. Patients and nurses determine variation in adherence to guidelines at Dutch hospitals more than internists or settings, 586

Dimock, J. see Shotliff, K.

Dina, C. see Lacquemant, C.

Dinh, S. see Kidron, M.

Dissanayake, A.M., Gamble, G., Baker, J.R. & Reed, L. Diagnostic delays of adult patients admitted to hospital with diabetic ketoacidosis, 290

Dobson, L., Sheldon, C.D. & Hattersley, A.T. Conventional measures underestimate glycaemia in cystic fibrosis patients, 691

Dodd, L. see O'Hare, J.P.

Doering, W. see Otter, W.

Dorn, J. see Menon, V.

Dotevall, A., Johansson, S., Wilhelmsen, L. & Rosengren, A. Increased levels of triglycerides, BMI and blood pressure and low physical activity increase the risk of diabetes in Swedish women. A prospective 18-year follow-up of the BEDAstudy, 615

Douglas, M. see Lawton, J.

Dubey, A.K. see Grant, R.W.

Duc Son, L.E.N.T., Kusama, K., Hung, N.T.K., Loan, T.T.H., Van Chuyen, N., Kunii, D., Sakai, T. & Yamamoto, S. Prevalence and risk factors for diabetes in Ho Chi Minh City, Vietnam, 371 Duengler, F. see Hadjadj, S. Dumskyj, M. see Usher, D.

Duncan, A. see Brown, A.

Duncan, C. see Brown, A.

Duncan, M.E. see Goldacre, M.J.

Dunger, D.B. see Schultz, C.J.

Durand-Zaleski, I. see Varroud-Vial, M.

Durrington, P.N. see Thomason, M.J.

Dwyer, T. see Sale, M.M.

Eastham, I. see Nocon, A.

Eaves, I. see Tait, K.F.

Ebrahim, S. see Lawlor, D.A.

Edwards, D.G. see Lajer, M.

Ek, J. see Lauenborg, J.

Elasha, H. see Bennett, S.M.A.

Eldor, R., Raz, I., Ben Yehuda, A. & Boulton, A.J.M. New and experimental approaches to treatment of diabetic foot ulcers: a comprehensive review of emerging treatment strategies, 1161

Elkeles, R.S., Feher, M.D., Flather, M.D., Godsland, I.F., Nugara, F., Richmond, W., Rubens, M.B. & Wang, D. for the PREDICT Study Group. The association of coronary calcium score and conventional cardiovascular risk factors in Type 2 diabetic subjects asymptomatic for coronary heart disease (The PREDICT Study), 1129

Ellis, J.D., Leese, G., McAlpine, R., Cole, A., MacEwen, C.J., Baines, P.S., Crombie, I. & Morris, A.D. for the DARTS/MEMO Collaboration. Prevalence of Diabetic Eye disease in Tayside, Scotland (P-DETS) Study: methodology, 1353

Emslie-Smith, A. see James, P.

Enoch, S. see Miller, D.R.

Eriksson, J.W. see Schölin, A. Eriksson, J.W. see Tan, M.H.

Erkens, J.A. see Spoelstra, J.A.

Ermini, G. see Marchesini, G.

Ernest, C.S. see Rave, K.

Esmatjes, E. see Flores, L.

Esteban, C. see Audí, L.

Eteiba, H. see Pell, J.P.

Eto, M. see Matsuda, M.

Evans, M.L., Hopkins, D., Macdonald, I.A. & Amiel, S.A. Alanine infusion during hypoglycaemia partly supports cognitive performance in healthy human subjects, 440

Faichney, A. see Pell, J.P.

Fanelli, C.G. see Porcellati, F.

Farooqui, K. see Jabbar, A.

Fattori, G. see Ciardullo, A.V.

Feher, M.D. see Al-Mrayat, M.

Feher, M.D. see Elkeles, R.S.

Feher, M.D. see Shotliff, K.

Feinglos, M.N. see Reboussin, D.M.

Feo, S. see Simmons, D.

Ferguson, S.C. see McAulay, V.

Fernández-Cancio, M. see Audí, L.

Ferrero, S. see Bruno, G.

Feskens, E.J.M. see van Dam, R.M.

Filioussi, K. see Bonovas, S.

Fineman, M. see Ratner, R.E.

Flapan, A. see Pell, J.P.



Flather, M.D. see Elkeles, R.S.

Fleckner, J. see Lajer, M.

Flores, L., Näf, S., Hernáez, R., Conget, I., Gomis, R. & Esmatjes, E. Transforming growth factor β_1 at clinical onset of Type 1 diabetes mellitus. A pilot study, 818

Flores, L., Vidal, M., Abian, J., Cases, A., Campistol, J.M., Clària, J., Lario, S. & Esmatjes, E. The effects of smoking and its cessation on 8-epi-PGF2α and transforming growth factor-beta 1 in Type 1 diabetes mellitus, 285

Fluiters, E. see Luna, R.

Foo, K., Sekhri, N., Knight, C., Deaner, A., Cooper, J., Ranjadayalan, K., Suliman, A. & Timmis, A.D. The effect of diabetes on heart rate and other determinants of myocardial oxygen demand in acute coronary syndromes, 1025

Foote, S. see Simmons, D.

Forlani, G. see Marchesini, G.

Formentini, G. see Bonora, E.

Forsén, A., Kangro, M., Sterner, G., Norrgren, K., Thorsson, O., Wollmer, P. & Sundkvist, G. A 14-year prospective study of autonomic nerve function in Type 1 diabetic patients: association with nephropathy, 852

Forst, T., Standl, E., Hohberg, C., Konrad, T., Schulze, J., Strotmann, H.J., Lübben, G., Pahler, S., Bachinger, A., Langenfeld, M. & Pfützner, A. IRIS II study: the IRIS II score assessment of a new clinical algorithm for the classification of insulin resistance in patients with Type 2 diabetes, 1149

Fou, F. see Simmons, D.

Franklyn, J.A. see Tait, K.F.

Fraser, R.J.L. see Russo, A.

Freemantle, N. see New, J.P.

Freudenheim, J.L. see Menon, V.

Friberg, H.H. see Heller, S.R.

Frier, B.M. see Graveling, A.J.

Frier, B.M. see McAulay, V.

Fritz-Hansen, T. see Taskiran, M.

Froguel, P. see Lacquemant, C.

Fujiwara, M., Baba, T., Neugebauer, S., Hasegawa, K., Hosoya, E., Tanaka, K., Shimada, K., Yamada, D. & Watanabe, T. Postprandial hypoglycaemia due to insulin lispro, 297

Fuller, J.H. see Thomason, M.J.

Fumelli, P. see Boemi, M.

Funatsu, H. see Mimura, T.

Furuhashi, M., Ura, N., Nakata, T., Tanaka, T. & Shimamoto, K. Genotype in human CD36 deficiency and diabetes mellitus, 952

Gale, E.A.M. & Atkinson, M. A piece of nucleic acid surrounded by controversy: coxsackievirus and the causes of Type 1 diabetes, 503

Galland-Jos, C. see Vergès, B.

Gallen, I. see Holt, R.I.G.

Gallery, E.D.M. see McGinn, S.

Galloway, T.S. see Devendra, D.

Gamble, G. see Dissanayake, A.M.

Gamble, G.D. see Simmons, D.

Game, F. see Gazis, A.

Game, F.L. see Jeffcoate, W.J.

Game, F.L. see Treece, K.A.

Garcia-Mayor, R.V. see Luna, R.

Gardener, E.A. see Khan, A.S.A.

Garon, J. see Gerstein, H.C.

Garuz, F. see Rigalleau, V.

Garvican, L. & Scanlon, P.H. A pilot quality assurance scheme for diabetic retinopathy risk reduction programmes, 1066

Gatling, W. see Swerdlow, A.J.

Gawron, W., Pospiech, L., Noczynska, A. & Koziorowska, E. Sudden hearing loss as a first complication of long-standing Type 1 diabetes mellitus: a case report, 96

Gazis, A., Pound, N., Macfarlane, R., Treece, K., Game, F. & Jeffcoate, W. Mortality in patients with diabetic neuropathic osteoarthropathy (Charcot foot), 1243

Geerlings, S.E. see Meiland, R.

Gemma, L. see Bonora, E.

George, A.M., Bodmer, C.W. & Aitken, J.M. Can the method of glycosylated haemoglobin assay affect results of such long-term studies?. 499

Gerstein, H.C., Garon, J., Joyce, C., Rolfe, A. & Walter, C.M.

Pre-prandial vs. post-prandial capillary glucose measurements as
targets for repaglinide dose titration in people with diet-treated or
metformin-treated Type 2 diabetes: a randomized controlled
clinical trial, 1200

van Gerwen, W.H.E.M. see Naushahi, M.J.

Getsios, D. see Caro, J.J.

Gibney, J., Turner, B., Weis, U., Meeking, D.R., Cansfield, J., Watts, G.F., Shaw, K.M. & Cummings, M.H. Reduced forearm reactive hyperaemia in normoalbuminuric subjects with Type 1 diabetes and retinopathy, 931

Gibson, J.M. see Khan, A.S.A.

Gibson, J.M. see New, J.P.

Gilbert, R.E. see Jerums, G.

Gilbey, S.G. see Rajeswaran, C.

Gildesgame, C. see Grant, R.W.

Gill-Randall, R.J., Adams, D., Ollerton, R.L. & Alcolado, J.C. Is human Type 2 diabetes maternally inherited? Insights from an animal model, 759

Gin, H. see Rigalleau, V.

Giunti, S. see Bruno, G.

Glümer, C. see Lauenborg, I.

Glümer, C., Jørgensen, T. & Borch-Johnsen, K. Targeted screening for undiagnosed diabetes reduces the number of diagnostic tests. Inter99(8), 874

Glover, M. see Moustafa, M.

Glueck, C.J., Bornovali, S., Pranikoff, J., Goldenberg, N., Dharashivkar, S. & Wang, P. Metformin, pre-eclampsia, and pregnancy outcomes in women with polycystic ovary syndrome, 829

Godsland, I.F. see Elkeles, R.S.

Goerzer, E. see Daxboeck, F.

Goff Jr, D.C. see Reboussin, D.M.

Goldacre, M.J., Duncan, M.E., Cook-Mozaffari, P. & Neil, H.A.W. Trends in mortality rates for death-certificate-coded diabetes mellitus in an English population 1979–99, 936

Goldberg, M. see Kidron, M.

Goldenberg, N. see Glueck, C.J.

Goldman, S.M. Diabetic peripheral neuropathy and spinal stenosis: prevalence of overlap and misdiagnosis. An introductory report,

Golmard, J.L. see Hartemann-Heurtier, A.

Gomis, R. see Flores, L.

Gonzales, C. see Rave, K.

Gooding, K. see Stride, A.

Gottsäter, A., Kangro, M. & Sundkvist, G. Early parasympathetic neuropathy associated with elevated fasting plasma C-peptide concentrations and late parasympathetic neuropathy with hyperglycaemia and other microvascular complications, 1304

Goudswaard, A.N., Stolk, R.P., Zuithoff, N.P.A., de Valk, H.W. & Rutten, G.E.H.M. Long-term effects of self-management education for patients with Type 2 diabetes taking maximal oral hypoglycaemic therapy: a randomized trial in primary care, 491

Gough, S.C.L. see Tait, K.F.

Gourgiotis, D. see Karagianni, C.

Grande, C. see Hillman, N.

Grant, R.W., Cagliero, E., Dubey, A.K., Gildesgame, C., Chueh, H.C., Barry, M.J., Singer, D.E., Nathan, D.M. & Meigs, J.B. Clinical inertia in the management of Type 2 diabetes metabolic risk factors, 150

Grassi, G. see Monge, L.

de Grauw, W.J.C. see Naushahi, M.J.

Graveling, A.J., Warren, R.E. & Frier, B.M. Hypoglycaemia and driving in people with insulin-treated diabetes: adherence to recommendations for avoidance, 1014

Grechi, G. see Mazzanti, L.

Green, B.N. see Reynolds, T.M.

Green, J., Casabonne, D. & Newton, R. Coxsackie B virus serology and Type 1 diabetes mellitus: a systematic review of published case-control studies, 507

Greenaway, T.M. see Sale, M.M.

Greenwood, R.G. see Winocour, P.H.

Gregory, I.W. see Lowes, L.

Gregson, C.L. & Allain, T.J. Amelanotic malignant melanoma disguised as a diabetic foot ulcer, 924

Griffin, S. see Whitford, D.L.

Griffiths, J.D. see McEwan, P.

Grill, V. see Qvigstad, E.

Grimaldi, A. see Hartemann-Heurtier, A.

Grimshaw, G. see Wilson, A.

Grobbee, D.E. see Spoelstra, J.A.

Grol, R.P.T.M. see Dijkstra, R.F.

Gross, J.L. see Costa, L.A.

Guilloteau, G. see Sallé, A.

Gul, A. see Jafar, T.H.

Gulliford, M. & Latinovic, R. Variations in glucose self-monitoring during oral hypoglycaemic therapy in primary care, 685

Gulliford, M.C., Mahabir, D. & Rocke, B. Diabetes-related inequalities in health status and financial barriers to health care access in a population-based study, 45

van Gurp, P.J.M. & Holleman, F. The 21st meeting of the Anglo-Danish-Dutch Diabetes Group (ADDDG), 99

Gussinyé, M. see Audí, L.

Gyntelberg, F. see Jørgensen, M.E.

Haak, T. see Kubiak, T.

Habib, A. see Jabbar, A.

Haddow, D. see Moustafa, M.

Hadjadj, S., Coisne, D., Mauco, G., Ragot, S., Duengler, F., Sosner, P., Torremocha, F., Herpin, D. & Marechaud, R. Prognostic value of admission plasma glucose and HbA_{1c} in acute myocardial infarction, 305

Hale, P.J. see Warburton, T.J.

Hale, P.M. see Raskin, P.

Hallevy, H. see Vander, T.

Hällsten, K., Virtanen, K.A., Lönnqvist, F., Janatuinen, T., Turiceanu, M., Rönnemaa, T., Viikari, J., Lehtimäki, T., Knuuti, J. & Nuutila, P. Enhancement of insulin-stimulated myocardial glucose uptake in patients with Type 2 diabetes treated with rosiglitazone, 1280

Halse, J. see Tan, M.H.

Hamilton, J. see Zdravkovic, V.

Hammond, J.J. see Jerums, G.

Han, K.H. see Kim, N.H.

Han, S.-M. see Kim, E.S.

Han, S.Y. see Kim, N.H.

Hanif, W. see O'Hare, J.P.

Hanley, A.J.G. see Retnakaran, R.

Hansen, B.V. see Lajer, M.

Hansen, T. see Lauenborg, J.

Haque, N. see Jabbar, A.

Harding, K.G. see Miller, D.R.

Hare, D.L. see Srivastava, P.M.

Harrison, S.R. see Nocon, A.

Hartemann-Heurtier, A., Robert, J., Jacqueminet, S., Ha Van, G., Golmard, J.L., Jarlier, V. & Grimaldi, A. Diabetic foot ulcer and multidrug-resistant organisms: risk factors and impact, 710

Harvey, J.N. & Allagoa, B. The long-term renal and retinal outcome of childhood-onset Type 1 diabetes, 26

Harvey, T.C. see Reynolds, T.M.

Hasegawa, K. see Fujiwara, M.

Hashimoto, K. see Ikeda, Y.

Hashimoto, K., Kasayama, S., Yamamoto, H., Kurebayashi, S., Kawase, I. & Koga, M. Strong association of C-reactive protein with body mass index and 2-h post-challenge glucose in non-diabetic, non-smoker subjects without hypertension, 581

Hassler-Hurst, J., Wadham, C. & Rayman, G. A double-blind study comparing 0.5% and 1% tropicamide for annual retinal screening in diabetic adolescents, 434

Hatcher, S. see Luzio, S.

Hattersley, A.T. see Dobson, L.

Hattersley, A.T. see Stride, A.

Ha Van, G. see Hartemann-Heurtier, A.

ten Have, P. see Dijkstra, R.F.

Hazelwood, K. see Sale, M.M.

Heine, R.J. see Adriaanse, M.C.

Heine, R.J. see DeVries, J.H.

Heine, R.J., Balkau, B., Ceriello, A., Del Prato, S., Horton, E.S. & Taskinen, M-R. What does postprandial hyperglycaemia mean?, 208

Heinemann, L. see Kapitza, C.

Heinemann, L. see Rave, K.

Heise, M. see Bennett, S.M.A.

Heise, T. see Kapitza, C.

Heldgaard, P.E., Olivarius, N. de. F., Hindsberger, C. & Henriksen, J.E. Impaired fasting glycaemia resembles impaired glucose tolerance with regard to cardiovascular risk factors: population-based, cross-sectional study of risk factors for cardiovascular disease,

Heller, S. Competing Interests, 961

Heller, S. Thoughts of a departing Editor, 1261

Heller, S. see Moustafa, M.

Heller, S. see Shearer, A.

Heller, S. & Kinmonth, A.L. A clinical research network in diabetes for the UK, 1061



Heller, S.R., Colagiuri, S., Vaaler, S., Wolffenbuttel, B.H.R., Koelendorf, K., Friberg, H.H., Windfeld, K. & Lindholm, A. Hypoglycaemia with insulin aspart: a double-blind, randomised, crossover trial in subjects with Type 1 diabetes, 769

Helminen, M. see Salminen, K.K. Henriksen, J.E. see Heldgaard, P.E.

Herishanu, Y.O. see Ifergane, G.

Herishanu, Y.O. see Vander, T.

Hermanns, N. see Kubiak, T.

Hernáez, R. see Flores, L.

Herpin, D. see Hadjadj, S.

Herranz, L. see Hillman, N.

Herrington, D.M. see Reboussin, D.M.

Herz, M. see Tan, M.H.

Hetyési, K. see Jermendy, G.

Heward, J.M. see Tait, K.F.

Hewer, W. see Mussell, M.

Hidvégi, T. see Jermendy, G.

Higa, M., Kaneko, Y. & Inokuchi, T. Two cases of hyperglycemic chorea in diabetic patients, 196

Higa, M., Takasu, N., Tamanaha, T., Nakamura, K., Shimabukuro, M., Sasara, T. & Tawata, M. Nitroglycerin spray rapidly improves pain in a patient with chronic painful diabetic neuropathy, 1053

Higa, N. see Shimabukuro, M.

Hiilesmaa, V. see Leinonen, A.

Hiilesmaa, V. see Loukovaara, S.

Hillhouse, E.W. see O'Hare, J.P.

Hillman, N., Herranz, L., Grande, C., Vaquero, P.M. & Pallardo, L.F. What is the relative contribution of blood glucose levels at different time points of the day to HbA_{1c} in Type 1 diabetes?, 468

Hillson, R.M. see Ng, R.S.H.

Hilsted, J. see Taskiran, M.

Himaga, M. see Usher, D.

Hindi, A. see Slater, R.A.

Hindsberger, C. see Heldgaard, P.E.

Hirai, C. see Seino, Y.

Hirst, J. Hypoglycaemia unawareness implicated: the value of

patient self-reporting, 194

Hitchman, R. see McElduff, A. Hitman, G.A. see Boucher, B.J.

Hitman, G.A. see Thomason, M.J.

Hod, M. see Ben-Haroush, A.

Hoebee, B. see van Dam, R.M.

Hoepelman, A.I.M. see Meiland, R.

Hoffmann, R. see Ortlepp, J.R.

Hogg, K.J. see Pell, J.P.

Hohberg, C. see Forst, T.

Holleman, F. see van Gurp, P.J.M.

Holman, C.D.J. see Bulsara, M.K.

Holman, R.R. see Cull, C.A.

Holt, R.I.G. & Coleman, M.A.G. Reply, 648

Holt, R.I.G. & Gallen, I. Time to move beyond glycosylated haemoglobin, 655

Holt, R.I.G., Peveler, R.C. & Byrne, C.D. Schizophrenia, the metabolic syndrome and diabetes, 515

Hopkins, D. see Evans, M.L.

Hopkinson, P. see McEwan, P.

Hori, S. see Mimura, T.

Hornnes, P. see Lauenborg, J.

Horowitz, M. see Russo, A.

Horton, E.S. see Heine, R.J.

Horton, S.J. see Devendra, D.

Horvath, K. see Siebenhofer, A.

Hosova, E. see Fujiwara, M. Howard, B.V. see Lu, W.Q.

Huh, K.B. see Ahn, C.W.

Huijsmans, Z. see Dijkstra, R.F.

Hummel, M., Ziegler, A.G. & Roth, R. Psychological impact of childhood islet autoantibody testing in families participating in the BABYDIAB study, 324

Hung, N.T.K. see Duc Son, L.E.N.T.

Hyöty, H. see Salminen, K.K.

Idris, I. see Jeffcoate, W.J.

Idris, I., Miller, D. & Page, S.R. Familial hyperinsulinaemia associated with epilepsy and mental retardation-a syndrome of familial insulin resistance, 628

Ifergane, G. see Vander, T.

Ikeda, Y. see Murata, M.

Ikeda, Y., Suehiro, T., Osaki, F., Tsuzura, S., Kumon, Y. & Hashimoto, K. Polymorphisms in the 5'-upstream region of the PKCB gene in Japanese patients with Type 2 diabetes, 1113

Ilonen, J. see Salminen, K.K.

Immonen, I. see Loukovaara, S.

Inokuchi, T. see Higa, M.

Inselmann, D. see Kubiak, T.

Irani, A. see Marita, A.R.

Irsigler, K. see Kästenbauer, T.

Ishibashi, S. see Nagasaka, S.

Ishida, K. see Ohshita, K.

Islam, N. see Jabbar, A.

Ito, T. see Nakashima, A.

Itoshima, T. see Nakatou, T.

Ivaldi, G. see Castelli, R.

Iwamoto, Y. see Mimura, T.

Izzo, P. see Lacquemant, C.

Jabbar, A., Farooqui, K., Habib, A., Islam, N., Haque, N. & Akhter, J. Clinical characteristics and outcomes of diabetic ketoacidosis in Pakistani adults with Type 2 diabetes mellitus, 920

Jablonski, K.A. see Lu, W.Q.

Jacqueminet, S. see Hartemann-Heurtier, A.

Jafar, T.H., Levey, A.S., White, F.M., Gul, A., Jessani, S., Khan, A.Q., Jafary, F.H., Schmid, C.H. & Chaturvedi, N. Ethnic differences and determinants of diabetes and central obesity among South Asians of Pakistan, 716

Jafary, F.H. see Jafar, T.H.

Jain, A.K. see Lu, W.Q.

James, P., Tan, H.H., MacAlpine, R., Brennan, G., Emslie-Smith, A. & Morris, A.D. on behalf of the DARTS/MEMO Collaboration. Treatment gap in the use of lipid-lowering drug therapy in diabetes: a population-based study, 1108

James, R.W. see Boemi, M.

Janatuinen, T. see Hällsten, K.

Jarlier, V. see Hartemann-Heurtier, A.

Jeffcoate, S.L. Diabetes control and complications: the role of glycated haemoglobin, 25 years on, 657

Jeffcoate, W. see Gazis, A.

Jeffcoate, W. The proper management of diabetic foot ulcers: time for a paradigm shift, 809

Jeffcoate, W.J. see Treece, K.A.

Jeffcoate, W.J., Idris, I. & Game, F.L. Erythromelia, or Mitchell's syndrome – new names for unexplained signs of inflammation in distal symmetrical neuropathy in diabetes, 1334

Jeffrey, R.R. see Pell, J.P.

Jennings, K. see Pell, J.P.

Jensen, G.B. see Taskiran, M.

Jermendy, G., Hetyési, K., Bíró, L. & Hidvégi, T. Prevalence of the metabolic syndrome in hypertensive and/or obese subjects, 805

Jerums, G. see Srivastava, P.M.

Jerums, G. see Thomas, M.C.

Jerums, G., Allen, T.J., Campbell, D.J., Cooper, M.E., Gilbert, R.E., Hammond, J.J., O'Brien, R.C., Raffaele, J. & Tsalamandris, C. on behalf of The Melbourne Diabetic Nephropathy Study Group. Long-term renoprotection by perindopril or nifedipine in non-hypertensive patients with Type 2 diabetes and microalbuminuria, 1192

Jessani, S. see Jafar, T.H.

Ji, Y.H. see Kim, N.H.

Jo, I. see Song, J.

Io, S.A. see Song, I.

Joglekar, A.A. see Kale, S.D.

Johannesen, J. see Larsen, Z.M.

Johansson, S. see Dotevall, A.

John, W.G. see Manley, S.

Johns, D. see Tan, M.H.

Johnson, J.A. see Yee, A.

Jones, A. see O'Hare, J.P.

Jones, K.L. see Lu, W.Q.

Jones, N.P. see Bennett, S.M.A.

Jones, T.W. see Bulsara, M.K.

Jorgensen, L.N. see Pan, C.Y.

Jorgensen, M.E., Bjerregaard, P., Gyntelberg, F. & Borch-Johnsen, K. and the Greenland Population Study. Prevalence of the metabolic syndrome among the Inuit in Greenland. A comparison between two proposed definitions, 1237

Jørgensen, M.E. & Borch-Johnsen, K. The metabolic syndrome—is one global definition possible?, 1064

Jørgensen, T. see Glümer, C.

Jørgensen, T. see Lauenborg, J.

Jovanovic, L. see Reboussin, D.M.

Joyce, C. see Gerstein, H.C.

Jung, M. see Song, J.

Jung, S.-C. see Song, J.

Kaaja, R. see Leinonen, A.

Kaaja, R.J. see Loukovaara, S.

Kaku, K. see Matsuda, M.

Kale, S.D., Yajnik, C.S., Kulkarni, S.R., Meenakumari, K., Joglekar, A.A., Khorsand, N., Ladkat, R.S., Ramdas, L.V. & Lubree, H.G. High risk of diabetes and metabolic syndrome in Indian women with gestational diabetes mellitus, 1257

Kamoshida, S. see Watanabe, A.

Kan, E.C.Y. see Ko, G.T.C.

Kanavakis, E. see Karagianni, C.

Kanda, Y. see Matsuda, M.

Kaneko, Y. see Higa, M.

Kang, Y.S. see Kim, N.H.

Kangro, M. see Forsén, A.

Kangro, M. see Gottsäter, A.

Kanou, M. see Watanabe, A.

Kapitza, C., Rave, K., Ostrowski, K., Heise, T. & Heinemann, L. Reduced postprandial glycaemic excursion with biphasic insulin Aspart 30 injected immediately before a meal, 500

Karagianni, C., Stabouli, S., Roumeliotou, K., Traeger-Synodinos, J., Kavazarakis, E., Gourgiotis, D., Lambrou, J. & Kanavakis, E. Severe hypertriglyceridaemia in diabetic ketoacidosis: clinical and genetic study, 380

Karasová, L. see Šimková, R.

Karlsson, F.A. see Schölin, A.

Karvonen, M. see Moltchanova, E.

Kasayama, S. see Hashimoto, K.

Kästenbauer, T., Sauseng, S., Brath, H., Abrahamian, H. & Irsigler, K. The value of the Rydel-Seiffer tuning fork as a predictor of diabetic polyneuropathy compared with a neurothesiometer, 563

Katoh, T. see Baba, T.

Kavazarakis, E. see Karagianni, C.

Kawasaki, F. see Matsuda, M.

Kawase, I. see Hashimoto, K.

Kaye, W. see Raskin, P.

Kazdová, L. see Šimková, R.

Kedar, Y. see Rachmani, R.

Keen, H. see Swerdlow, A.J.

Kempler, P. see Ziegler, D.

Kempler, P. Textbook of Diabetic Neuropathy, 400

Kerner, A. see Aronson, D.

Kerr, D. see Chevne, E.H.

Khalid, B.A.K. see Pan, C.Y.

Khan, A.Q. see Jafar, T.H.

Khan, A.S.A., Talbot, J.A., Tieszen, K.L., Gardener, E.A., Gibson, J.M. & New, J.P. Evaluation of a bedside blood ketone sensor: the effects of acidosis, hyperglycaemia and acetoacetate on sensor performance, 782

Khorsand, N. see Kale, S.D.

Khunti, K.K. see Davies, M.J.

Khutoryansky, N. see Raskin, P.

Kidron, M., Dinh, S., Menachem, Y., Abbas, R., Variano, B., Goldberg, M., Arbit, E. & Bar-On, H. A novel per-oral insulin formulation: proof of concept study in non-diabetic subjects, 354

Kim, C.S., Kim, D.M., Park, J.S., Nam, J.Y., Ahn, C.W., Cha, B.S., Lim, S.K., Kim, K.R. & Lee, H.C. A case of primary antiphospholipid syndrome and Type 2 diabetes mellitus with large artery thromboses successfully treated by abdominal stent implantation, 1258

Kim, D.L. see Kim, N.H.

Kim, D.M. see Ahn, C.W.

Kim, D.M. see Kim, C.S.

Kim, E. see Song, J.

Kim, E.S., Han, S.-M., Kim, Y.I., Song, K.-H., Kim, M.-S., Kim, W.B., Park, J.-Y. & Lee, K.-U. Prevalence and clinical characteristics of metabolic syndrome in a rural population of South Korea, 1141

Kim, K.B. see Kim, N.H.

Kim, K.R. see Ahn, C.W.

Kim, K.R. see Kim, C.S.

Kim, M.-S. see Kim, E.S.

Kim, N.H., Kim, K.B., Kim, D.L., Kim, S.G., Choi, K.M., Baik, S.H., Choi, D.S., Kang, Y.S., Han, S.Y., Han, K.H., Ji, Y.H. & Cha, D.R. Plasma and urinary vascular endothelial growth factor and diabetic nephropathy in Type 2 diabetes mellitus, 545

Kim, S.G. see Kim, N.H.

Kim, S.-S. see Song, J.

Kim, W.B. see Kim, E.S.



Kim, Y.I. see Kim, E.S.

Kinjo, Y. see Shirayama, H.

Kinmonth, A.L. see Heller, S.

Kitano, S. see Mimura, T.

Kleybrink, S. see Otter, W.

Klittich, W.S. see Caro, J.J.

Klungel, O.H. see Spoelstra, J.A.

Knight, C. see Foo, K.

Knip, M. see Salminen, K.K.

Knuuti, J. see Hällsten, K.

Ko, G.T.C., Li, J.K.Y., Kan, E.C.Y. & Lo, M.K.W. Effects of a structured health education programme by a diabetic education nurse on cardiovascular risk factors in Chinese

Type 2 diabetic patients: a 1-year prospective randomized study, 1274

Kockum, I. see Schölin, A.

Koelendorf, K. see Heller, S.R.

Koga, M. see Hashimoto, K.

Kogure, A. see Yoshioka, K.

Kohner, E. see Luzio, S.

Kohno, N. see Ohshita, K.

Koistinen, R. see Loukovaara, S.

Kollind, M. see Qvigstad, E.

Kolterman, O.G. see Ratner, R.E.

Komine, F. see Watanabe, A.

Kondo, K. see Seino, Y.

Kong, M.-F. see Russo, A.

Konkoy, C.S. see Tan, M.H.

Konrad, T. see Forst, T.

Kotsopoulos, D. see Meyer, C.

Kousa, A. see Moltchanova, E.

Koziorowska, E. see Gawron, W.

Krause, R. see Daxboeck, F.

Kristiansen, O.P. see Larsen, Z.M.

Kristiansen, O.P. see Myrup, B.

Krsek, M. see Silha, J.V.

Kubiak, T., Hermanns, N., Schreckling, H.J., Kulzer, B. & Haak, T. Assessment of hypoglycaemia awareness using continuous glucose monitoring, 487

Kubiak, T., Kuhr, B., Inselmann, D., Hermanns, N., Kulzer, B. & Haak, T. Reversible cognitive deterioration after a single episode of severe hypoglycaemia: a case report, 1366

Kuhr, B. see Kubiak, T.

Kulkarni, S.R. see Kale, S.D.

Kulzer, B. see Kubiak, T.

Kulzer, B. see Mussell, M.

Kumar, S. see O'Hare, J.P.

Kumar, S. see Valsamakis, G.

Kumon, Y. see Ikeda, Y.

Kunii, D. see Duc Son, L.E.N.T.

Kurebayashi, S. see Hashimoto, K.

Kurukulasuriya, N. see Spallone, V.

Kusama, K. see Duc Son, L.E.N.T.

Laakso, M. see Sivenius, K.

Laatikainen, L. see Loukovaara, S.

Lacquemant, C., Froguel, P., Lobbens, S., Izzo, P., Dina, C. & Ruiz, J. The adiponectin gene SNP+45 is associated with coronary artery disease in Type 2 (non-insulin-dependent) diabetes mellitus, 776 Ladkat, R.S. see Kale, S.D.

Laing, S.P. see Swerdlow, A.J.

Lajer, M., Tarnow, L., Fleckner, J., Hansen, B.V., Edwards, D.G., Parving, H-H. & Boel, E. Association of aldose reductase gene Z+2 polymorphism with reduced susceptibility to diabetic nephropathy in Caucasian Type 1 diabetic patients, 867

Lambrou, J. see Karagianni, C.

Landin-Olsson, M. see Schölin, A.

Langenfeld, M. see Forst, T.

Lario, S. see Flores, L.

Larsen, Z.M., Johannesen, J., Kristiansen, O.P., Nerup, J. & Pociot, F.,
The Danish IDDM Epidemiology and Genetics Group, The
Danish Study Group of IDDM in Childhood. Evidence for
linkage on chromosome 4p16.1 in Type 1 diabetes Danish
families and complete mutation scanning of the WFS1
(Wolframin) gene, 218

Larsson, H.B.W. see Taskiran, M.

Latinovic, R. see Gulliford, M.

Lauenborg, J., Damm, P., Ek, J., Glümer, C., Jørgensen, T., Borch-Johnsen, K., Vestergaard, H., Hornnes, P., Pedersen, O. & Hansen, T. Studies of the Ala/Val98 polymorphism of the hepatocyte nuclear factor-1 α gene and the relationship to β -cell function during an OGTT in glucose-tolerant women with and without previous gestational diabetes mellitus, 1310

Lawlor, D.A., Ebrahim, S. & Smith, G.D. The metabolic syndrome and coronary heart disease in older women: findings from the British Women's Heart and Health Study, 906

Lawton, J., Peel, E., Douglas, M. & Parry, O. 'Urine testing is a waste of time': newly diagnosed Type 2 diabetes patients' perceptions of self-monitoring, 1045

Lazarovitch, T. see Slater, R.A.

Leakehe, L. see Simmons, D.

Lean, M.E.J. see Bates, C.J.

Lee, H.C. see Ahn, C.W.

Lee, H.C. see Kim, C.S.

Lee, H.K. see Song, J.

Lee, J.H. see Ahn, C.W.

Lee, K.-U. see Kim, E.S.

Lee, P.P. see Palinkas, L.A.

Lee, W-J. see Yu, H-I.

Leese, G. see Ellis, J.D.

Lehtimäki, T. see Hällsten, K.

Leinonen, A., Hiilesmaa, V., Andersen, H., Teramo, K. & Kaaja, R. Diurnal blood glucose profiles in women with gestational diabetes with or without hypertension, 1181

Lemieux, S. see Piché, M.E.

Leone, A. see Wang, X.L.

Leppert, K. see Braun, A.

Letondeur, C. see Varroud-Vial, M.

Leufkens, H.G.M. see Spoelstra, J.A.

Levey, A.S. see Jafar, T.H.

Levi, Z. see Rachmani, R.

Levy, J.C. see Wallace, T.M.

Levy, Y. see Aronson, D.

Li Voon Chong, J.S.W. see Teh, M.M.

Li, J.K.Y. see Ko, G.T.C.

Liesenfeld, B. see Luzio, S.

Lim, H.S. & Lip, G.Y.H. Impaired fasting glycaemia vs. impaired glucose tolerance: two sides of the same coin with implications for cardiovascular risk, 303 Lim, S.K. see Ahn, C.W.

Lim, S.K. see Kim, C.S.

Lindholm, A. see Heller, S.R.

Lip, G.Y.H. see Lim, H.S.

Lipkin, E.W. see Reboussin, D.M.

Lisbôa, H.R.K. see Costa, L.A.

van de Lisdonk, E.H. see Naushahi, M.I.

Liu, H-C. see Yu, H-I.

Livingstone, S.J. see Thomason, M.J.

Lo Presti, R. see Caimi, G.

Lo, M.K.W. see Ko, G.T.C.

Loan, T.T.H. see Duc Son, L.E.N.T.

Lobbens, S. see Lacquemant, C.

Lombardi, S. see Bonora, E.

Lönnqvist, F. see Hällsten, K.

Loukovaara, S., Immonen, I., Koistinen, R., Rudge, J., Teramo, K.A., Laatikainen, L., Hiilesmaa, V. & Kaaja, R.J. Angiopoietic factors and retinopathy in pregnancies complicated with Type 1 diabetes, 697

Lovecchio, M. see Spallone, V.

Low, P.A. see Ziegler, D.

Lowes, L., Lyne, P. & Gregory, J.W. Childhood diabetes: parents' experience of home management and the first year following diagnosis, 531

Lu, W.Q., Resnick, H.E., Jablonski, K.A., Jain, A.K., Jones, K.L., Robbins, D.C. & Howard, B.V. Effects of glycaemic control on cardiovascular disease in diabetic American Indians: the Strong Heart Study, 311

Lübben, G. see Forst, T.

Lubree, H.G. see Kale, S.D.

Lucas, B. see Vergès, B.

Ludvigsson, J. see Samuelsson, U.

Luna, R., Fluiters, E., Rodriguez, I. & Garcia-Mayor, R.V. Growth in Type 1 diabetic children, 1054

Lunt, M.J. see Cheyne, E.H.

Lutale, J.K. see Morbach, S.

Luzio, S., Hatcher, S., Zahlmann, G., Mazik, L., Morgan, M., Liesenfeld, B., Bek, T., Schuell, H., Schneider, S., Owens, D.R. & Kohner, E. Feasibility of using the TOSCA telescreening procedures for diabetic retinopathy, 1121

Lyne, P. see Lowes, L.

MacAlpine, R. see James, P.

Macdonald, I.A. see Evans, M.L.

MacEwen, C.I. see Ellis, I.D.

MacFarlane, I.A. see Daousi, C.

Macfarlane, I.A. see Saunders, S.A.

Macfarlane, R. see Gazis, A.

Macfarlane, R.M. see Treece, K.A.

MacIsaac, R.J. see Srivastava, P.M.

MacIsaac, R.J. see Thomas, M.C.

Mackness, M.I. see Thomason, M.J.

MacNeil, S. see Moustafa, M.

McAlister, F.A. see Yee, A.

McAlpine, R. see Ellis, J.D.

McAulay, V., Ferguson, S.C. & Frier, B.M. Post-prandial administration of insulin lispro with a high fat meal minimizes risk of hypoglycaemia in Type 1 diabetes, 953

McBain, A.M. see Brown, A.

McCallum, R.W. see McEleavy, O.D.

McElduff, A. & Hitchman, R. Fasting plasma glucose values alone miss most abnormalities of glucose tolerance in the postpartum, 648

McEleavy, O.D., McCallum, R.W., Petrie, J.R., Small, M.,

Connell, J.M.C., Sattar, N. & Cleland, S.J. Higher carotid-radial pulse wave velocity in healthy offspring of patients with Type 2 diabetes, 262

McEwan, P., Williams, J.E., Griffiths, J.D., Bagust, A., Peters, J.R., Hopkinson, P. & Currie, C.J. Evaluating the performance of the Framingham risk equations in a population with diabetes, 318

McGill, J. see Raskin, P.

McGinn, S., Poronnik, P., Gallery, E.D.M. & Pollock, C.A. The effects of high glucose and atorvastatin on endothelial cell matrix production, 1102

McGrath, B.P. see Meyer, C.

McIntosh, C. see Al-Mrayat, M.

McKee, M. see Shapo, L.

Ma, R.C.W. see Chan, W.B.

Madsbad, S. see Tan, M.H.

Maggs, D.G. see Ratner, R.E.

Maghenzani, G. see Monge, L. Mahabir, D. see Gulliford, M.C.

Mahaney, M.C. see Wang, X.L.

Maiello, M.R. see Spallone, V.

Majumdar, S.R. see Yee, A. Manini, R. see Marchesini, G.

Mankad, P. see Pell, J.P.

Manley, S., John, W.G. & Marshall, S. Introduction of IFCC reference method for calibration of HbA_{1c}: implications for clinical care, 673

Mansoor, M.A. see Bates, C.J.

Maranian, M. see Tait, K.F.

Marchesini, G., Forlani, G., Cerrelli, F., Manini, R., Natale, S., Baraldi, L., Ermini, G., Savorani, G., Zocchi, D. & Melchionda, N. WHO and ATPIII proposals for the definition of the metabolic syndrome in patients with Type 2 diabetes, 383

Marechaud, R. see Hadjadj, S.

Marini, F. see Bonora, E.

Marita, A.R., Rane, S., Mokal, R.A., Nair, S.R. & Irani, A. Autoantibodies against GAD65 and IA-2 in recently diagnosed Type 1 diabetic children from Western India, 956

Markiewicz, W. see Aronson, D.

Marra, G. see Ceriello, A.

Marshall, A.L., Chetwynd, A., Morris, J.A., Placzek, M., Smith, C., Olabi, A. & Thistlethwaite, D. Type 1 diabetes mellitus in childhood: a matched case control study in Lancashire and Cumbria, UK, 1035

Marshall, S. see Manley, S.

Marshall, S.M. Inhibition of the renin–angiotensin system: added value in reducing cardiovascular and renal risk?, 1

Marshall, T. see Tait, K.F.

Martí, G. see Audí, L.

Martin, J. see Saunders, S.A.

Martinelli, L. see Ceriello, A.

Maruyama, T. see Murata, M.

Masaki, T. see Nakashima, A.

Mason, J.M. see New, J.P.

Matsuda, M., Kawasaki, F., Yamada, K., Kanda, Y., Saito, M., Eto, M., Matsuki, M. & Kaku, K. Impact of adiposity and plasma adipocytokines on diabetic angiopathies in Japanese Type 2 diabetic subjects, 881



Matsuki, M. see Matsuda, M.

Matthews, D.R. see Wallace, T.M.

Mauco, G. see Hadjadj, S.

Mazik, L. see Luzio, S.

Mazzanti, L., Nanetti, L., Vignini, A., Rabini, R.A., Grechi, G., Cester, N., Curzi, C.M. & Tranquilli, A.L. Gestational diabetes affects platelet behaviour through modified oxidative radical metabolism, 68

Meddah, D. see Cosson, E.

Meeking, D.R. see Gibney, J.

Meenakumari, K. see Kale, S.D.

Meigs, J.B. see Grant, R.W.

Meiland, R., Geerlings, S.E., De Neeling, A.J. & Hoepelman, A.I.M.
Diabetes mellitus in itself is not a risk factor for antibiotic resistance
in *Escherichia coli* isolated from patients with bacteriuria, 1032

Melchionda, N. see Marchesini, G.

Menachem, Y. see Kidron, M.

Mennuni, G. see Spallone, V.

Menon, V., Ram, M., Dorn, J., Armstrong, D., Muti, P., Freudenheim, J.L., Browne, R., Schunemann, H. & Trevisan, M. Oxidative stress and glucose levels in a population-based sample, 1346

Menzinger, G. see Spallone, V.

Mercadé, I. see Rodríguez-Villar, C.

Metrikat, J. see Ortlepp, J.R.

Meyer, C., Milat, F., McGrath, B.P., Cameron, J., Kotsopoulos, D. & Teede, H.J. Vascular dysfunction and autonomic neuropathy in Type 2 diabetes, 746

Milat, F. see Meyer, C.

Miller, D. see Idris, I.

Miller, D.R., Enoch, S., Blow, M., Harding, K.G. & Price, P.E.

Effectiveness of a new brand of stock 'diabetic' shoes to protect
against diabetic foot ulcer relapse. A prospective cohort study, 646

Mimura, T., Funatsu, H., Uchigata, Y., Kitano, S., Shimizu, E., Noma, H., Amano, S., Araie, M., Iwamoto, Y. & Hori, S. Development and progression of diabetic retinopathy in patients with Type 1 diabetes who are positive for GAD autoantibody, 559

Mishra, K. see O'Hare, J.P.

Mitamura, Y. see Tashimo, A.

Mizue, Y. see Tashimo, A.

Mogensen, C.E. & Cooper, M.E. Diabetic renal disease: from recent studies to improved clinical practice, 4

Mohan, V. see Pan, C.Y.

Mokal, R.A. see Marita, A.R.

Möllenberg, J. see Morbach, S.

Møller, F. see Sjølie, A.K.

Moltchanova, E., Rytkönen, M., Kousa, A., Taskinen, O., Tuomilehto, J. & Karvonen, M. for the Spat Study Group, the Finnish Childhood Diabetes Registry Group. Zinc and nitrate in the ground water and the incidence of Type 1 diabetes in Finland, 256

Monge, L., Bruno, G., Pinach, S., Grassi, G., Maghenzani, G., Dani, F. & Pagano, G. A clinically orientated approach increases the efficiency of screening for latent autoimmune diabetes in adults (LADA) in a large clinic-based cohort of patients with diabetes onset over 50 years, 456

Montgomery, S.M. Reply, 398

Moore, D. see Shotliff, K.

Morbach, S., Lutale, J.K., Viswanathan, V., Möllenberg, J., Ochs, H.R., Rajashekar, S., Ramachandran, A. & Abbas, Z.G. Regional differences in risk factors and clinical presentation of diabetic foot lesions, 91 Morgan, M. see Luzio, S.

Mori, M. see Shimizu, H.

Morris, A.D. see Ellis, J.D.

Morris, A.D. see James, P.

Morris, A.D. see Swerdlow, A.J.

Morris, J.A. see Marshall, A.L.

Motz, E. see Ceriello, A.

Moustafa, M., Simpson, C., Glover, M., Dawson, R.A., Tesfaye, S., Creagh, F.M., Haddow, D., Short, R., Heller, S. & Macneil, S. A new autologous keratinocyte dressing treatment for non-healing diabetic neuropathic foot ulcers, 786

Muchmore, D. see Rave, K.

Muggeo, M. see Bonora, E.

Mughal, S. see O'Hare, J.P.

Muller, R. see Braun, A.

Muller, U.A. see Braun, A.

Munro, N. see Al-Mrayat, M.

Murata, M., Maruyama, T., Suzuki, Y., Saruta, T. & Ikeda, Y.
Paraoxonase 1 ¹⁹²Gln/Arg polymorphism is associated with the risk of microangiopathy in Type 2 diabetes mellitus, 837

Murday, A. see Pell, J.P.

Murphy, L.J. see Nyomba, B.L.G.

Murphy, L.J. see Silha, J.V.

Mussell, M., Hewer, W., Kulzer, B., Bergis, K. & Rist, F. Effects of improved glycaemic control maintained for 3 months on cognitive function in patients with Type 2 diabetes, 1253

Muti, P. see Menon, V.

Myrup, B., Yokoyama, H., Kristiansen, O.P., Østergaard, P.B. & Olivecrona, T. Release of endothelium-associated proteins into blood by injection of heparin in normal subjects and in patients with Type 1 diabetes, 1135

Nabe, K. see Watanabe, A.

Nadeau, A. see Piché, M.E.

Näf, S. see Flores, L.

Nagai, S. see Tashimo, A.

Nagasaka, S., Aiso, Y., Yoshizawa, K. & Ishibashi, S. Comparison of pioglitazone and metformin efficacy using homeostasis model assessment, 136

Nair, S.R. see Marita, A.R.

Nakachi, K. see Shirayama, H.

Nakamura, A. see Nakatou, T.

Nakamura, K. see Higa, M.

Nakamura, S. see Watanabe, A.

Nakamura, Y. see Tashimo, A.

Nakanishi, N., Suzuki, K. & Tatara, K. Haematocrit and risk of development of Type 2 diabetes mellitus in middle-aged Japanese men, 476

Nakashima, A., Nakashima, R., Ito, T., Masaki, T. & Yorioka, N. HMG-CoA reductase inhibitors prevent bone loss in patients with Type 2 diabetes mellitus, 1020

Nakashima, R. see Nakashima, A.

Nakata, K. see Nakatou, T.

Nakata, T. see Furuhashi, M.

Nakatou, T., Nakata, K., Nakamura, A. & Itoshima, T. Carotid haemodynamic parameters as risk factors for cerebral infarction in Type 2 diabetic patients, 223

Nam, J.H. see Ahn, C.W.

Nam, J.Y. see Kim, C.S.

Nanetti, L. see Mazzanti, L.

Natale, S. see Marchesini, G.

Nathan, D.M. see Grant, R.W.

Naushahi, M.J., de Grauw, W.J.C., Avery, A.J., van Gerwen, W.H.E.M., van de Lisdonk, E.H. & van Weel, C. Risk factors for development of impaired renal function in Type 2 diabetes mellitus patients in primary care, 1096

Negishi, M. see Shimizu, H.

Nehr, M. see Daxboeck, F.

Neil, H.A.W. see Cull, C.A.

Neil, H.A.W. see Goldacre, M.J.

Neil, H.A.W. see Schultz, C.J.

Neil, H.A.W. see Thomason, M.J.

Nerup, J. see Larsen, Z.M.

Nesto, R. C-reactive protein, its role in inflammation, Type 2 diabetes and cardiovascular disease, and the effects of insulin-sensitizing treatment with thiazolidinediones, 810

Neugebauer, S. see Fujiwara, M.

Neugebauer-Baba, S. see Baba, T.

New, J.P. see Khan, A.S.A.

New, J.P., Mason, J.M., Freemantle, N., Teasdale, S., Wong, L., Bruce, N.J., Burns, J.A. & Gibson, J.M. Educational outreach in diabetes to encourage practice nurses to use primary care hypertension and hyperlipidaemia guidelines (EDEN): a randomized controlled trial, 599

Newton, R. see Green, J.

Ng, M.C.Y. see Chan, W.B.

Ng, R.S.H., Darko, D.A. & Hillson, R.M. Street drug use among young patients with Type 1 diabetes in the UK, 295

Nichols, G.A. & Brown, J.B. Functional status before and after diagnosis of Type 2 diabetes, 793

Nielsen, J.F., Andersen, H. & Sinkjær, T. Decreased stiffness at the ankle joint in patients with long-term Type 1 diabetes, 539

Niipels, G. see Adriaanse, M.C.

Nirei, K. see Watanabe, A.

Nishihira, J. see Tashimo, A.

Niskanen, L. see Sivenius, K.

Nobels, F., Beckers, F., Bailleul, E., De Schrijver, P., Sierens, L. & Van Crombrugge, P. Feasibility of a quality assurance programme of bedside blood glucose testing in a hospital setting: 7 years' experience, 1288

Nocon, A., Rhodes, P.J., Wright, J.P., Eastham, J., Williams, D.R.R., Harrison, S.R. & Young, R.J. Specialist general practitioners and diabetes clinics in primary care: a qualitative and descriptive evaluation, 32

Noczynska, A. see Gawron, W.

Noma, H. see Mimura, T.

Norrgren, K. see Forsén, A.

Nosek, L. see Rave, K.

Nowak, H. see Ziegler, D.

Nozawa, Y. see Baba, T.

Nugara, F. see Elkeles, R.S.

Nurmikko, T.J. see Daousi, C.

Nussey, S. see Usher, D.

Nuutila, P. see Hällsten, K.

Nyomba, B.L.G. see Silha, J.V.

Nyomba, B.L.G., Berard, L. & Murphy, L.J. Facilitating access to glucometer reagents increases blood glucose self-monitoring frequency and improves glycaemic control: a prospective study in insulin-treated diabetic patients, 129

Nyström, L. see Schölin, A.

O'Brien, J.A. see Caro, J.J.

O'Brien, R.C. see Jerums, G.

Ochs, H.R. see Morbach, S.

O'Hare, J.P., Raymond, N.T., Mughal, S., Dodd, L., Hanif, W., Ahmad, Y., Mishra, K., Jones, A., Kumar, S., Szczepura, A., Hillhouse, E.W. & Barnett, A.H. on behalf of the UKADS study group. Evaluation of delivery of enhanced diabetes care to patients of South Asian ethnicity: the United Kingdom Asian Diabetes Study (UKADS), 1357

Oh-I, S. see Shimizu, H.

Ohshiro, Y. see Shirayama, H.

Ohshita, K., Yamane, K., Ishida, K., Watanabe, H., Okubo, M. & Kohno, N. Post-challenge hyperglycaemia is an independent risk factor for arterial stiffness in Japanese men, 636

Ohtsuka, K. see Tashimo, A.

Oikarinen, S. see Salminen, K.K.

Okazaki, Y. see Seino, Y.

Okubo, H. see Watanabe, A.

Okubo, M. see Ohshita, K.

Okumura, A. see Seino, Y.

Olabi, A. see Marshall, A.L.

Oldroyd, K. see Pell, J.P.

Oli, J.M., Watkins, P.J., Wild, B. & Adegoke, O.J. Albuminuria in Afro-Caribbeans with Type 2 diabetes mellitus: is the sickle cell trait a risk factor?, 483

Olivarius, N...F. see Heldgaard, P.E.

Olivecrona, T. see Myrup, B.

Ollerton, R.L. see Gill-Randall, R.J.

Ortlepp, J.R., Metrikat, J. & Hoffmann, R. Reply, 803

Osaki, F. see Ikeda, Y.

Osaki, R. see Chan, W.B.

Østerby, R. see Baba, T.

Østergaard, P.B. see Myrup, B.

Östman, J. see Schölin, A.

Ostrowski, K. see Kapitza, C.

Ota, T. see Tsuchiyama, N.

Otsuka, M. see Watanabe, A.

Otter, W., Kleybrink, S., Doering, W., Standl, E. & Schnell, O. Hospital outcome of acute myocardial infarction in patients with and without diabetes mellitus, 183

Owens, D.R. see Luzio, S.

Oyarzabal, M. see Audí, L.

Oyibo, S.O., Breislin, K. & Boulton, A.J.M. Electrical stimulation therapy through stocking electrodes for painful diabetic neuropathy: a double blind, controlled crossover study, 940

Pagano, G. see Bruno, G.

Pagano, G. see Monge, L.

Page, S.R. see Idris, I.

Pahler, S. see Forst, T.

Palinkas, L.A., Lee, P.P. & Barrett-Connor, E. A prospective study of Type 2 diabetes and depressive symptoms in the elderly: The Rancho Bernardo Study, 1185

Pallardo, L.F. see Hillman, N.

Palmer, A.J., Roze, S., Valentine, W.J. & Spinas, G. The costeffectiveness of continuous subcutaneous insulin infusion compared with multiple daily injections for the management of diabetes: response to Schuffman and Carr, 1372

Pampanelli, S. see Porcellati, F.



Pan, C.Y., So, W.Y., Khalid, B.A.K., Mohan, V., Thai, A.C., Zimmet, P., Cockram, C.S., Jorgensen, L.N. & Yeo, J.P., The ASDIAB Study Group. Metabolic, immunological and clinical characteristics in newly diagnosed Asian diabetes patients aged 12-40 years, 1007

Panahloo, A. see Wood, D.M.

Paries, J. see Cosson, E.

Park, J.S. see Kim, C.S.

Park, J.-Y. see Kim, E.S.

Park, S.W. see Ahn, C.W.

Parry, O. see Lawton, J.

Parving, H-H. see Lajer, M.

Pascot, A. see Piché, M.E.

Patois-Vergès, B. see Vergès, B.

Patterson, C.C. see Swerdlow, A.J.

Patterson, D.J., Thirlby, R.C. & Sobrino, M.A. High-frequency gastric stimulation in a patient with diabetic gastroparesis, 195

Paycha, F. see Cosson, E.

Pearson, E.R. see Stride, A.

Pedersen, O. see Lauenborg, J.

Peel, E. see Lawton, J.

Pelikánová, T. see Šimková, R.

Pell, A.C.H. see Pell, J.P.

Pell, J.P., Pell, A.C.H., Jeffrey, R.R., Jennings, K., Oldroyd, K., Eteiba, H., Hogg, K.J., Murday, A., Faichney, A., Colquhoun, I., Berg, G., Starkey, I.R., Flapan, A. & Mankad, P. Comparison of survival following coronary artery bypass grafting vs. percutaneous coronary intervention in diabetic and non-diabetic patients: retrospective cohort study of 6320 procedures, 790

Peponis, V. see Bonovas, S.

Perbellini, S. see Bonora, E.

Pérez-Heras, A. see Rodríguez-Villar, C.

Perin, P.C. see Bruno, G.

Perriello, G. see Porcellati, F.

Peters, J.R. see McEwan, P.

Peters, S. see Dijkstra, R.F.

Petit, C. see Varroud-Vial, M.

Petrie, J.R. see McEleavy, O.D.

Pettitt, D.J. see Reboussin, D.M.

Peveler, R.C. see Holt, R.I.G.

Pfützner, A. see Forst, T.

Piché, M.E., Després, J.P., Pascot, A., Nadeau, A., Tremblay, A., Weisnagel, S.J., Bergeron, J. & Lemieux, S. Impaired fasting glucose vs. glucose intolerance in pre-menopausal women: distinct metabolic entities and cardiovascular disease risk?, 730

Piconi, L. see Ceriello, A.

Pieber, T.R. see Siebenhofer, A.

Pinach, S. see Monge, L.

Placzek, M. see Marshall, A.L.

Plank, J. see Siebenhofer, A.

van der Ploeg, H.M. see Adriaanse, M.C.

Pociot, F. see Larsen, Z.M.

Poli, M. see Bonora, E.

Pollock, C.A. see Mcginn, S.

Porcellati, F., Rossetti, P., Pampanelli, S., Fanelli, C.G., Torlone, E., Scionti, L., Perriello, G. & Bolli, G.B. Better long-term glycaemic control with the basal insulin glargine as compared with NPH in patients with Type 1 diabetes mellitus given meal-time lispro insulin, 1213

Poronnik, P. see Mcginn, S.

Porro, T. see Castelli, R.

Porta, M. & Trento, M. on behalf of the ROMEO Writing Committee. ROMEO: rethink organization to improve education and outcomes, 644

Pospiech, L. see Gawron, W.

Pound, N. see Gazis, A.

Pound, N. see Treece, K.A.

Pranikoff, J. see Glueck, C.J.

Pratley, R.E. see Stefan, N.

Prentice, A. see Bates, C.J.

Price, P.E. see Miller, D.R.

Probstfield, J.L. see Reboussin, D.M.

Qiao, Z. see Swerdlow, A.J.

Quagliaro, L. see Ceriello, A.

Qvigstad, E., Kollind, M. & Grill, V. Nine weeks of bedtime diazoxide is well tolerated and improves \beta-cell function in subjects with Type 2 diabetes, 73

Rabini, R.A. see Mazzanti, L.

Rachmani, R., Slavachevsky, I., Amit, M., Levi, Z., Kedar, Y., Berla, M. & Ravid, M. The effect of spironolactone, cilazapril and their combination on albuminuria in patients with hypertension and diabetic nephropathy is independent of blood pressure reduction: a randomized controlled study, 471

Raffaele, J. see Jerums, G.

Raffaelli, A. see Bonora, E.

Ragot, S. see Hadjadj, S.

Raif, N. see Retnakaran, R.

Rainwater, D.L. see Wang, X.L.

Rajashekar, S. see Morbach, S.

Rajeswaran, C. & Dr Gilbey, S.G. Oxford Handbook of Endocrinology and Diabetes, 199

Ram, M. see Menon, V.

Ramachandran, A. see Morbach, S.

Ramachandran, A. see Viswanathan, V.

Ramadan, A. see Cosson, E.

Ramdas, L.V. see Kale, S.D.

Ramot, Y. see Slater, R.A.

Rane, S. see Marita, A.R.

Ranjadayalan, K. see Foo, K.

Rapoport, M.J. see Slater, R.A.

Raskin, P., McGill, J., Saad, M.F., Cappleman, J.M., Kaye, W., Khutoryansky, N. & Hale, P.M. for the Repaglinide/ Rosiglitazone Study Group. Combination therapy for Type 2 diabetes: repaglinide plus rosiglitazone, 329

Rasmussen, B. see Taskiran, M.

Rasmussen, V. see Taskiran, M.

Ratner, R.E., Dickey, R., Fineman, M., Maggs, D.G., Shen, L., Strobel, S.A., Weyer, C. & Kolterman, O.G. Amylin replacement with pramlintide as an adjunct to insulin therapy improves long-term glycaemic and weight control in Type 1 diabetes mellitus: a 1-year, randomized controlled trial, 1204

Rave, K. see Kapitza, C.

Rave, K., Nosek, L., Heinemann, L., Gonzales, C., Ernest, C.S., Chien, J. & Muchmore, D. Inhaled micronized crystalline human insulin using a dry powder inhaler: dose-response and time-action profiles, 763

Ravid, M. see Rachmani, R.

Rayman, G. see Hassler-Hurst, J.

Raymond, N.T. see O'Hare, J.P.

Raz, I. see Eldor, R.

Reboussin, D.M., Goff Jr, D.C., Lipkin, E.W., Herrington, D.M., Summerson, J., Steffes, M., Crouse III, R.J., Jovanovic, L., Feinglos, M.N., Probstfield, J.L., Banerji, M.A., Pettitt, D.J. & Williamson, J. The combination oral and nutritional treatment of late-onset diabetes mellitus (CONTROL DM) trial results, 1082

Reed, L. see Dissanayake, A.M.

Resnick, H.E. see Lu, W.Q.

Retnakaran, R., Hanley, A.J.G., Raif, N., Connelly, P.W., Sermer, M. & Zinman, B. Hypoadiponectinaemia in South Asian women during pregnancy: evidence of ethnic variation in adiponectin concentration. 388

Reynolds, T.M., Twomey, P.J., Harvey, T.C. & Green, B.N. The number of unexpected HbA_{1c} variants may be a greater problem in routine practice than is generally realized, 1041

Rhodes, P.J. see Nocon, A.

Richmond, W. see Elkeles, R.S.

Rigalleau, V., Barbe, M., Garuz, F., Baillet-Blanco, L., Beauvieux, M.C. & Gin, H. Nutritional factors and diabetic foot: a role for vitamin F2-951

Rist, F. see Mussell, M.

Ritz, P. see Sallé, A.

Robbins, D.C. see Lu, W.Q.

Robert, J. see Hartemann-Heurtier, A.

Roberts, S. see Shearer, A.

Roberts, S.H. see Whitford, D.L.

Rocke, B. see Gulliford, M.C.

Rodriguez, I. see Luna, R.

Rodríguez-Villar, C., Pérez-Heras, A., Mercadé, I., Casals, E. & Ros, E.
Comparison of a high-carbohydrate and a high-monounsaturated
fat, olive oil-rich diet on the susceptibility of LDL to oxidative
modification in subjects with Type 2 diabetes mellitus, 142

Rolfe, A. see Gerstein, H.C.

Rönnemaa, T. see Hällsten, K.

Ros, E. see Rodríguez-Villar, C.

Rosengren, A. see Dotevall, A.

Rossetti, P. see Porcellati, F.

Roth, R. see Hummel, M.

Roumeliotou, K. see Karagianni, C.

Rowe, B. see Tait, K.F.

Roze, S. see Palmer, A.J.

Rubens, M.B. see Elkeles, R.S.

Rudge, J. see Loukovaara, S.

Ruiz, J. see Lacquemant, C.

Russo, A., Botten, R., Kong, M.-F., Chapman, I.M., Fraser, R.J.L., Horowitz, M. & Sun, W.-M. Effects of acute hyperglycaemia on anorectal motor and sensory function in diabetes mellitus, 176

Rutten, G.E.H.M. see Goudswaard, A.N.

Rutten, G.E.H.M. see Spoelstra, J.A.

Ryan, M. see Sallé, A.

Rytkönen, M. see Moltchanova, E.

Saad, M.F. see Raskin, P.

Saggiani, F. see Bonora, E.

Saito, M. see Matsuda, M.

Sakai, T. see Duc Son, L.E.N.T.

Sakurai, K. see Baba, T.

Sakurai, M. see Tsuchiyama, N.

Sale, M.M., Hazelwood, K., Zimmet, P.Z., Shaw, J.E., Stankovich, J.M., Greenaway, T.M. & Dwyer, T. Trends in diabetes management practices of patients from an Australian insulin-treated diabetes register. 165

Sallé, A., Guilloteau, G., Ryan, M., Bouhanick, B. & Ritz, P. Effect of insulin treatment on the body composition of Type 2 diabetic patients, 1298

Salminen, K.K., Vuorinen, T., Oikarinen, S., Helminen, M., Simell, S., Knip, M., Ilonen, J., Simell, O. & Hyöty, H. Isolation of enterovirus strains from children with preclinical Type 1 diabetes, 156

Samarasinghe, Y. see Al-Mrayat, M.

Samuelsson, U. & Ludvigsson, J. The concentrations of short-chain fatty acids and other microflora-associated characteristics in faeces from children with newly diagnosed Type 1 diabetes and control children and their family members, 64

Sanderson, D. see Shearer, A.

Santi, L. see Bonora, E.

dos Santos Silva, I. see Swerdlow, A.J.

Saruta, T. see Murata, M.

Sasara, T. see Higa, M.

Sattar, N. see McEleavy, O.D.

Saunders, S.A., Democratis, J., Martin, J. & Macfarlane, I.A. Intravenous drug abuse and Type 1 diabetes: financial and healthcare implications, 1269

Sauseng, S. see Kästenbauer, T.

Savorani, G. see Marchesini, G.

Sawada, S. see Watanabe, A.

Sawcer, S. see Tait, K.F.

Scanlon, P.H. see Garvican, L.

Scarpitta, A.M. see Caimi, G.

Schaap, M.M. see van Dam, R.M.

Schernthaner, G., Wein, W., Shnawa, N., Bates, P.C. & Birkett, M.A. Preprandial vs. postprandial insulin lispro—a comparative crossover trial in patients with Type 1 diabetes, 279

Schiel, R. see Braun, A.

Schmid, C.H. see Jafar, T.H.

Schneider, S. see Luzio, S.

Schnell, O. see Otter, W.

Schölin, A., Törn, C., Nyström, L., Berne, C., Arnqvist, H., Blohmé, G., Bolinder, J., Eriksson, J.W., Kockum, I., Landin-Olsson, M., Östman, J., Karlsson, F.A., Sundkvist, G. & Björk, E. Normal weight promotes remission and low number of islet antibodies prolong the duration of remission in Type 1 diabetes, 447

Schreckling, H.J. see Kubiak, T.

Schuell, H. see Luzio, S.

Schultz, C.J., Dalton, R.N., Selwood, M., Dunger, D.B. & Neil, H.A.W. on behalf of the Oxford Regional Prospective Study Group. Paternal phenotype is associated with microalbuminuria in young adults with Type 1 diabetes mellitus of short duration, 246

Schulze, J. see Forst, T.

Schunemann, H. see Menon, V.

Scionti, L. see Porcellati, F.

Scuffham, P.A. & Carr, L. Reply, 1372

Seena, R. see Viswanathan, V.

Seidell, J.C. see van Dam, R.M.

Seino, Y., Yamauchi, M., Hirai, C., Okumura, A., Kondo, K., Yamamoto, M. & Okazaki, Y. A case of fulminant Type 1 diabetes associated with mexiletine hypersensitivity syndrome, 1156

Sekhri, N. see Foo, K.

Selwood, M. see Schultz, C.J.



Sermer, M. see Retnakaran, R.

Shapo, L., McKee, M., Coker, R. & Ylli, A. Type 2 diabetes in Tirana City, Albania: a rapid increase in a country in transition, 77

Shaw, J. see Delaney, C.

Shaw, J. see Zimmet, P.

Shaw, J.E. see Sale, M.M.

Shaw, K. see Winocour, P.H.

Shaw, K.M. see Gibney, J.

Shearer, A., Bagust, A., Sanderson, D., Heller, S. & Roberts, S. Cost-effectiveness of flexible intensive insulin management to enable dietary freedom in people with Type 1 diabetes in the UK, 460

Sheldon, C.D. see Dobson, L.

Shen, L. see Ratner, R.E.

Sherwin, R.S. see Cheyne, E.H.

Sheu, W.H-H. see Yu, H-I.

Shimabukuro, M. see Higa, M.

Shimabukuro, M., Higa, N., Takasu, N., Tagawa, T. & Ueda, S. A single dose of nateglinide improves post-challenge glucose metabolism and endothelial dysfunction in Type 2 diabetic patients, 983

Shimada, K. see Fujiwara, M.

Shimamoto, K. see Furuhashi, M.

Shimizu, E. see Mimura, T.

Shimizu, H., Shimomura, K., Negishi, M., Oh-I, S., Tomita, Y., Uehara, Y. & Mori, M. Glycaemic control and the increase in pulse wave velocity (PWV) in Type 2 diabetic patients, 804

Shimomura, K. see Shimizu, H.

Shin, C. see Song, J.

Shirayama, H., Ohshiro, Y., Kinjo, Y., Taira, S., Teruya, I., Nakachi, K., Tawata, M. & Takasu, N. Acute brain injury in hypoglycaemia-induced hemiplegia, 623

Shitman, E. see Aronson, D.

Shnawa, N. see Schernthaner, G.

Short, R. see Moustafa, M.

Shotliff, K. see Al-Mrayat, M.

Shotliff, K., Moore, D., Dimock, J. & Feher, M.D. Screening for diabetic retinopathy—false positives do occur (it could be Shagreene), 651

Siebenhofer, A., Plank, J., Horvath, K., Berghold, A., Sutton, A.J., Sommer, R. & Pieber, T.R. Angiotensin receptor blockers as anti-hypertensive treatment for patients with diabetes mellitus: meta-analysis of controlled double-blind randomized trials, 18

Sierens, L. see Nobels, F.

Silha, J.V., Krsek, M., Skrha, J., Sucharda, P., Nyomba, B.L.G. & Murphy, L.J. Plasma resistin, leptin and adiponectin levels in non-diabetic and diabetic obese subjects, 497

Simek, S. see Šimková, R.

Simell, O. see Salminen, K.K.

Simell, S. see Salminen, K.K.

Šimková, R., Kazdová, L., Karasová, L., Simek, S. & Pelikánová, T.

Effect of acute hyperglycaemia on sodium handling and excretion
of nitric oxide metabolites, bradykinin, and cGMP in Type 1
diabetes mellitus, 968

Simmons, D., Gamble, G.D., Foote, S., Cole, D.R. & Coster, G. The New Zealand Diabetes Passport Study: a randomized controlled trial of the impact of a diabetes passport on risk factors for diabetes-related complications, 214

Simmons, D., Voyle, J.A., Fou, F., Feo, S. & Leakehe, L. Tale of two churches: differential impact of a church-based diabetes control programme among Pacific Islands people in New Zealand, 122 Simon, D. see Varroud-Vial, M.

Simpson, C. see Moustafa, M.

Simpson, S.H. see Yee, A.

Sinagra, D. see Caimi, G.

Singer, D.E. see Grant, R.W.

Sinkjær, T. see Nielsen, J.F.

Sirolla, C. see Boemi, M.

Sivagami, M. see Viswanathan, V.

Sivenius, K., Niskanen, L., Voutilainen-Kaunisto, R., Laakso, M. & Uusitupa, M. Aldose reductase gene polymorphisms and susceptibility to microvascular complications in Type 2 diabetes, 1325

Sjølie, A.K. & Møller, F. Medical management of diabetic retinopathy, 666

Skrha, J. see Silha, J.V.

Slater, R.A., Lazarovitch, T., Boldur, I., Ramot, Y., Buchs, A., Weiss, M., Hindi, A. & Rapoport, M.J. Swab cultures accurately identify bacterial pathogens in diabetic foot wounds not involving hone. 705

Slater, S.D. see Swerdlow, A.J.

Slavachevsky, I. see Rachmani, R.

Small, M. see McEleavy, O.D.

Smith, C. see Marshall, A.L.

Smith, G.D. see Lawlor, D.A.

Snehalatha, C. see Viswanathan, V.

Snoek, F.J. see Adriaanse, M.C.

Snoek, F.J. see DeVries, J.H.

Snook, H. see Tait, K.F.

So, W.Y. see Chan, W.B.

So, W.Y. see Pan, C.Y. Sobrino, M.A. see Patterson, D.J.

Sommer, R. see Siebenhofer, A.

Song, J., Kim, E., Shin, C., Kim, S.-S., Lee, H.K., Jung, M., Jung, S.-C., Jo, S.A. & Jo, I. Prevalence of the metabolic syndrome among South Korean adults: the Ansan study, 1154

Song, K.-H. see Kim, E.S.

Song, S.H. & Brown, P.M. Coronary heart disease risk assessment in diabetes mellitus: comparison of UKPDS risk engine with Framingham risk assessment function and its clinical implications, 238

Song, Y.D. see Ahn, C.W.

Song, Y-M. see Yu, H-I.

Sosner, P. see Hadjadj, S.

Spallone, V., Maiello, M.R., Kurukulasuriya, N., Barini, A., Lovecchio, M., Tartaglione, R., Mennuni, G. & Menzinger, G. Does autonomic neuropathy play a role in erythropoietin regulation in non-proteinuric Type 2 diabetic patients?, 1174

Spijkerman, A.M.W. see Adriaanse, M.C.

Spinas, G. see Palmer, A.J.

Spoelstra, J.A., Stolk, R.P., Klungel, O.H., Erkens, J.A., Rutten, G.E.H.M., Leufkens, H.G.M. & Grobbee, D.E. Initiation of glucose-lowering therapy in Type 2 diabetes mellitus patients in general practice, 896

Srivastava, P.M., Calafiore, P., Macisaac, R.J., Hare, D.L., Jerums, G. & Burrell, L.M. Thiazolidinediones and congestive heart failure—exacerbation or new onset of left ventricular dysfunction?, 945

Stabouli, S. see Karagianni, C.

Standl, E. see Forst, T.

Standl, E. see Otter, W.

Stankovich, J.M. see Sale, M.M.

Starkey, I.R. see Pell, J.P.

Steel, J.M. see Brown, A.

Stefan, N., Stumvoll, M., Weyer, C., Bogardus, C., Tataranni, P.A. & Pratley, R.E. Exaggerated insulin secretion in Pima Indians and African-Americans but higher insulin resistance in Pima Indians compared to African-Americans and Caucasians, 1090

Stefanadis, C. see Antoniades, C.

Steffes, M. see Reboussin, D.M.

Sterner, G. see Forsén, A.

Stolk, R.P. see Goudswaard, A.N.

Stolk, R.P. see Spoelstra, J.A.

Strand, J. see Tan, M.H.

Strauss, K. see Varroud-Vial, M.

Stride, A., Pearson, E.R., Brown, A., Gooding, K., Castleden, H.A.J. & Hattersley, A.T. Serum amino acids in patients with mutations in the hepatocyte nuclear factor-1 alpha gene, 928

Strobel, S.A. see Ratner, R.E.

Strotmann, H.J. see Forst, T.

Stumvoll, M. see Stefan, N.

Sucharda, P. see Silha, J.V.

Suehiro, T. see Ikeda, Y.

Suliman, A. see Foo, K.

Summerson, J. see Reboussin, D.M.

Sun, W.-M. see Russo, A.

Sundkvist, G. see Forsén, A.

Sundkvist, G. see Gottsäter, A.

Sundkvist, G. see Schölin, A.

Sutton, A.J. see Siebenhofer, A.

Suzuki, K. see Nakanishi, N.

Suzuki, Y. see Murata, M.

Swerdlow, A.J., Laing, S.P., dos Santos Silva, L., Slater, S.D., Burden, A.C., Botha, J.L., Waugh, N.R., Morris, A.D., Gatling, W., Bingley, P.J., Patterson, C.C., Qiao, Z. & Keen, H. Mortality of South Asian patients with insulin-treated diabetes mellitus in the United Kingdom: a cohort study, 845

Szczepura, A. see O'Hare, J.P.

Tagawa, T. see Shimabukuro, M.

Taira, S. see Shirayama, H.

Tait, K.F., Collins, J.E., Heward, J.M., Eaves, I., Snook, H., Franklyn, J.A., Barnett, A.H., Todd, J.A., Maranian, M., Compston, A., Sawcer, S. & Gough, S.C.L. Evidence for a Type 1 diabetes-specific mechanism for the insulin gene-associated IDDM2 locus rather than a general influence on autoimmunity, 267

Tait, K.F., Marshall, T., Berman, J., Carr-Smith, J., Rowe, B., Todd, J.A., Bain, S.C., Barnett, A.H., Gough, S.C.L. Clustering of autoimmune disease in parents of siblings from the Type 1 diabetes Warren repository, 358

Takakura, Y. see Yoshioka, K.

Takamura, T. see Tsuchiyama, N.

Takasu, N. see Higa, M.

Takasu, N. see Shimabukuro, M.

Takasu, N. see Shiravama, H.

Talbot, J.A. see Khan, A.S.A.

Tamanaha, T. see Higa, M.

Tamura, K. see Watanabe, A.

Tan, H.H. see James, P.

Tan, M.H., Johns, D., Strand, J., Halse, J., Madsbad, S., Eriksson, J.W., Clausen, J., Konkoy, C.S. & Herz, M. for the GLAC Study Group. Sustained effects of pioglitazone vs. glibenclamide on insulin sensitivity, glycaemic control, and lipid profiles in patients with Type 2 diabetes, 859

Tanaka, K. see Fujiwara, M.

Tanaka, T. see Furuhashi, M.

Targher, G. see Bonora, E.

Tarnow, L. see Lajer, M.

Tartaglione, R. see Spallone, V.

Tashimo, A., Mitamura, Y., Nagai, S., Nakamura, Y., Ohtsuka, K., Mizue, Y. & Nishihira, J. Aqueous levels of macrophage migration inhibitory factor and monocyte chemotactic protein-1 in patients with diabetic retinopathy, 1292

Tasker, P. Management of Type II Diabetes in Primary Care: A Practical Guide, 958

Taskinen, M-R. see Heine, R.J.

Taskinen, O. see Moltchanova, E.

Taskiran, M., Rasmussen, V., Rasmussen, B., Fritz-Hansen, T., Larsson, H.B.W., Jensen, G.B. & Hilsted, J. Left ventricular dysfunction in normotensive Type 1 diabetic patients: the impact of autonomic neuropathy, 524

Tatara, K. see Nakanishi, N.

Tataranni, P.A. see Stefan, N.

Tawata, M. see Higa, M.

Tawata, M. see Shirayama, H.

Taylor, R. see Dashora, U.K.

Teasdale, S. see New, J.P.

Teede, H.J. see Meyer, C.

Teh, M.M., Brooks, A.P. & Li Voon Chong, J.S.W. Metabolic targets achieved in elderly diabetic patients attending a secondary care annual review diabetic clinic, 1052

Tempesta, A. see Castelli, R.

Tentolouris, C. see Antoniades, C.

Teramo, K. see Leinonen, A.

Teramo, K.A. see Loukovaara, S.

Teruya, I. see Shirayama, H.

Tesfaye, S. see Moustafa, M. Testa, R. see Boemi, M.

Thai, A.C. see Pan, C.Y.

The DAI Study Group. The prevalence of coronary heart disease in Type 2 diabetic patients in Italy: the DAI study, 738

The Danish IDDM Epidemiology and Genetics Group and The Danish Study Group of IDDM in Childhood see Larsen, Z.M.

Thirlby, R.C. see Patterson, D.J.

Thistlethwaite, D. see Marshall, A.L.

Thomas, M.C., MacIsaac, R.J., Tsalamandris, C. & Jerums, G. Elevated iron indices in patients with diabetes, 798

Thomas, P.W. see Cheyne, E.H.

Thomas, S. see Coll, A.P.

Thomason, M.J., Colhoun, H.M., Livingstone, S.J., Mackness, M.I., Betteridge, D.J., Durrington, P.N., Hitman, G.A., Neil, H.A.W. & Fuller, J.H., the CARDS Investigators. Baseline characteristics in the Collaborative AtoRvastatin Diabetes Study (CARDS) in patients with Type 2 diabetes, 901

Thompson, J. see Wilson, A.

Thorsson, O. see Forsén, A.

Tieszen, K.L. see Khan, A.S.A.

Timmis, A.D. see Foo, K.

Toda, H. see Yoshioka, K.

Todd, J.A. see Tait, K.F.

Tomita, Y. see Shimizu, H.

Tong, P.C.Y. see Chan, W.B.

Torlone, E. see Porcellati, F.

Törn, C. see Schölin, A.

Torremocha, F. see Hadjadj, S.

Tountas, C. see Antoniades, C.

Tousoulis, D. see Antoniades, C. Toutouza, M. see Antoniades, C.

Toutouzas, P. see Antoniades, C.

Traeger-Synodinos, J. see Karagianni, C.

Tranquilli, A.L. see Mazzanti, L.

Treece, K. see Gazis, A.

Treece, K.A., Macfarlane, R.M., Pound, N., Game, F.L. & Jeffcoate, W.J. Validation of a system of foot ulcer classification

in diabetes mellitus, 987 Tremblay, A. see Piché, M.E.

Treml, H. see Al-Mravat, M.

Trento, M. see Porta, M.

Tres, G.S. see Costa, L.A.

Trevisan, M. see Menon, V.

Tringham, J.R. see Davies, M.J.

Troughton, J. see Davies, M.J.

Tsakok, A.D. The statistical analysis of cardiovascular risks, 955

Tsalamandris, C. see Jerums, G.

Tsalamandris, C. see Thomas, M.C.

Tseng, C.-H. Independent association of uric acid levels with peripheral arterial disease in Taiwanese patients with Type 2 diabetes, 724

Tsioufis, C. see Antoniades, C.

Tsuchiyama, N., Ando, H., Ota, T., Sakurai, M. & Takamura, T. Modulating effects of olanzapine on the development of diabetic ketoacidosis, 300

Tsuyuki, R.T. see Yee, A.

Tsuzura, S. see Ikeda, Y.

Tuomilehto, J. see Moltchanova, E.

Turiceanu, M. see Hällsten, K.

Turner, B. see Gibney, J.

Twisk, J.W.R. see Adriaanse, M.C.

Twomey, P.J. see Reynolds, T.M.

Uchigata, Y. see Mimura, T.

Uchiyama, T. see Watanabe, A.

Ueda, S. see Shimabukuro, M.

Uehara, Y. see Shimizu, H.

Umekawa, T. see Yoshioka, K.

Ura, N. see Furuhashi, M.

Usher, D., Dumskyj, M., Himaga, M., Williamson, T.H., Nussey, S. & Boyce, J. Automated detection of diabetic retinopathy in digital retinal images: a tool for diabetic retinopathy screening, 84

Uusitupa, M. see Sivenius, K.

Vaaler, S. see Heller, S.R.

Valensi, P. see Cosson, E.

Valentine, W.J. see Palmer, A.J.

de Valk, H.W. see Goudswaard, A.N.

Valsamakis, G., Chetty, R., Anwar, A., Banerjee, A.K., Barnett, A. & Kumar, S. Association of simple anthropometric measures of obesity with visceral fat and the metabolic syndrome in male Caucasian and Indo-Asian subjects, 1339

Van Chuyen, N. see Duc Son, L.E.N.T.

Van Crombrugge, P. see Nobels, F.

Vander, T., Hallevy, H., Ifergane, G. & Herishanu, Y.O.

Metformin-induced encephalopathy without lactic acidosis, 194

Vaquero, P.M. see Hillman, N.

Vargha, P. see Ziegler, D.

Variano, B. see Kidron, M.

Varroud-Vial, M., Simon, D., Attali, J., Durand-Zaleski, I., Bera, L., Attali, C., Letondeur, C., Strauss, K., Petit, C. & Charpentier, G. Improving glycaemic control of patients with Type 2 diabetes in a primary care setting: a French application of the Staged Diabetes Management programme, 592

Vasiliadou, C. see Antoniades, C.

Vergès, B., Patois-Vergès, B., Cohen, M., Lucas, B., Galland-Jos, C. & Casillas, J.M. Effects of cardiac rehabilitation on exercise capacity in Type 2 diabetic patients with coronary artery disease, 289

Vestergaard, H. see Lauenborg, J.

Veves, A. see Viswanathan, V.

Vidal, M. see Flores, L.

Vignini, A. see Mazzanti, L.

Viikari, J. see Hällsten, K.

Virtanen, K.A. see Hällsten, K.

Viswanathan, V. see Morbach, S. Viswanathan, V., Sivagami, M., Seena, R., Snehalatha, C.,

Ramachandran, A. & Veves, A. Increased forefoot to rearfoot plantar pressure ratio in South Indian patients with diabetic foot ulceration, 396

Voutilainen-Kaunisto, R. see Sivenius, K.

Voyle, J.A. see Simmons, D.

Vuorinen, T. see Salminen, K.K.

Wadham, C. see Hassler-Hurst, J.

Walker, M. see Bennett, S.M.A.

Wallace, T.M., Levy, J.C. & Matthews, D.R. An increase in insulin sensitivity and basal beta-cell function in diabetic subjects treated with pioglitazone in a placebo-controlled randomized study, 568

Walter, C.M. see Gerstein, H.C.

Wang, D. see Elkeles, R.S.

Wang, P. see Glueck, C.J.

Wang, X.L., Rainwater, D.L., Leone, A. & Mahaney, M.C. Effects of diabetes on plasma nitrotyrosine levels, 577

Warburton, T.J., Hale, P.J. & Dewhurst, J.A. Evaluation of a local optometric diabetic retinopathy screening service, 632

Warren, R.E. see Graveling, A.J.

Watanabe, A., Komine, F., Nirei, K., Tamura, K., Nabe, K., Aiba, N., Kamoshida, S., Otsuka, M., Okubo, H., Kanou, M., Sawada, S., Uchiyama, T., Nakamura, S. & Arakawa, Y. A case of secondary diabetes mellitus with acromegaly improved by pioglitazone, 1049

Watanabe, H. see Ohshita, K.

Watanabe, T. see Baba, T.

Watanabe, T. see Fujiwara, M.

Watkins, P.J. see Oli, J.M.

Watts, G.F. see Gibney, J.

Waugh, N.R. see Swerdlow, A.J.

van Weel, C. see Naushahi, M.J. Wein, W. see Schernthaner, G.

Weis, U. see Gibney, J.

Weisnagel, S.J. see Piché, M.E.

Weiss, M. see Slater, R.A.

Weyer, C. see Ratner, R.E.

Weyer, C. see Stefan, N.

White, F.M. see Jafar, T.H.

Whitford, D.L. & Roberts, S.H. Changes in prevalence and site of care of diabetes in a health district 1991–2001, 640

Whitford, D.L., Roberts, S.H. & Griffin, S. Sustainability and effectiveness of comprehensive diabetes care to a district population, 1221

Wild, B. see Oli, J.M.

Wilding, J.P.H. see Bennett, S.M.A.

Wilhelmsen, L. see Dotevall, A.

Wilkin, T.J. see Devendra, D.

Williams, D.R.R. see Nocon, A.

Williams, J.E. see McEwan, P.

Williamson, J. see Reboussin, D.M.

Williamson, T.H. see Usher, D.

Wilson, A., Baker, R., Thompson, J. & Grimshaw, G. Coverage in screening for diabetic retinopathy according to screening provision: results from a national survey in England and Wales, 271

Windfeld, K. see Heller, S.R.

Winocour, P.H., Shaw, K. & Greenwood, R.G. Access to dietetic services in the UK, 646

Wolffenbuttel, B.H.R. see Heller, S.R.

Wollmer, P. see Forsén, A.

Wong, L. see New, J.P.

Woo, S.O. see Ahn, C.W.

Wood, D.M., Athwal, S. & Panahloo, A. The advantages and disadvantages of a 'herbal' medicine in a patient with diabetes mellitus: a case report, 62.5

Woodward, A. see Daousi, C.

Wright, J.P. see Nocon, A.

Yajnik, C.S. see Kale, S.D.

Yamada, D. see Fujiwara, M.

Yamada, K. see Matsuda, M.

Yamamoto, H. see Hashimoto, K.

Yamamoto, M. see Seino, Y.

Yamamoto, S. see Duc Son, L.E.N.T.

Yamane, K. see Ohshita, K.

Yamauchi, M. see Seino, Y.

Yee, A., Majumdar, S.R., Simpson, S.H., Mcalister, F.A., Tsuyuki, R.T. & Johnson, J.A. Statin use in Type 2 diabetes mellitus is associated with a delay in starting insulin, 962

Yeo, J.P. see Pan, C.Y.

Yeste, D. see Audí, L.

Ylli, A. see Shapo, L.

Yogev, Y. see Ben-Haroush, A.

Yokoyama, H. see Myrup, B.

Yorioka, N. see Nakashima, A.

Yoshida, T. see Yoshioka, K.

Yoshikawa, T. see Yoshioka, K.

Yoshioka, K., Yoshida, T., Takakura, Y., Umekawa, T., Kogure, A., Toda, H. & Yoshikawa, T. Adiponectin gene polymorphism (G276T) and diabetic retinopathy in Japanese patients with Type 2 diabetes, 1158

Yoshioka, K., Yoshida, T., Umekawa, T., Kogure, A., Takakura, Y., Toda, H. & Yoshikawa, T. Methylenetetrahydrofolate reductase gene polymorphism is not related to diabetic nephropathy in Japanese Type 2 diabetic patients, 1051

Yoshizawa, K. see Nagasaka, S.

Young, R.J. see Nocon, A.

Yu, H-I., Sheu, W.H-H., Song, Y-M., Liu, H-C., Lee, W-J. & Chen, Y-T. C-reactive protein and risk factors for peripheral vascular disease in subjects with Type 2 diabetes mellitus, 336

Zahlmann, G. see Luzio, S.

Zdravkovic, V., Daneman, D. & Hamilton, J. Presentation and course of Type 2 diabetes in youth in a large multi-ethnic city, 1144

Zenari, L. see Bonora, E.

Ziegler, A.G. see Hummel, M.

Ziegler, D., Nowak, H., Kempler, P., Vargha, P. & Low, P.A. Treatment of symptomatic diabetic polyneuropathy with the antioxidant α -lipoic acid: a meta-analysis, 114

Zimmet, P. see Pan, C.Y.

Zimmet, P., Shaw, J. & Alberti, K.G.M.M. Reply, 650

Zimmet, P.Z. see Sale, M.M.

Zinder, O. see Aronson, D.

Zinman, B. see Retnakaran, R.

Zocchi, D. see Marchesini, G.

Zuithoff, N.P.A. see Goudswaard, A.N.

Subject Index

α-lipoic acid, 114

β-cell function, 188

β-cell hormone, 1204

B-cell rest, 73

β-cell, 1310

16S rDNA sequencing, 1247

75-g oral glucose tolerance test, 636

A_{1c}, 311

Aboriginal health, 188

access to health care, 45

access to nealth car acetoacetate, 782 acromegaly pioglitazone, 1049

acute coronary syndromes, 1025

acute myocardial infarction, 305

acute phase response, 203

adhesion molecules, 552

adipocytokines, 881

adiponectin, 388, 776

adolescent, 434, 1144

Afro-Caribbeans, 483

alanine, 440

Albania, 77

albuminuria, 471, 483

alcohol, 230

aldose reductase, 867, 1325

alternative medicine, 625

ambulatory blood pressure, 415

American Indians, 311

amphetamine, 295

amputation, 987

amylin, 1204

anaemia, 798, 1174 analysis, 1288

angioplasty, 790

angiopoietin-1, 697

angiopoietin-2, 697

angiotensin receptor blockers, 18

ankle joint, 539

ankle-brachial index, 336



anorectal motility, 176 antibiotic resistance, 1032 antibody screening, 324 anti-glutamic acid decarboxylase antibodies, 456 anti-inflammatory drugs, 666 anti-islet cell antibodies, 456 antioxidants, 1346 APM1 gene, 776 aqueous humour, 1292 arterial compliance, 746 arterial stiffness, 262, 636, 746 Asians, 1007 aspirin, 1368 atherosclerosis, 223 atorvastatin, 901, 1102 ATP III criteria, 1141 atypical anti-psychotic drugs, 515 autoantibodies, 1007 autoantibodies to glutamic acid decarboxylase, 559 autoimmune disease, 358 autoimmunity, 267 autonomic function, 746 autonomic neuropathy, 524, 852, 1174,

bacteriuria, 1032 basal insulin, 1213 bed time insulin, 73 beta-cell function, 568 bioelectrical impedance, 1298 bisphosphonates, 1334 blockers of the renin-angiotensin system, 666 blood glucose, 1288 blood glucose meter, 1045 blood glucose self-monitoring, 129, 685 blood pressure, 1025 BMI, 447, 615 body composition changes, 1298 body mass index, 349, 581 bone mineral density, 1020 Bordetella trematum, 1247 brachial artery ultrasound, 1082 bradykinin, 969 brain metabolism, 440 British, 677

Camperdown haemoglobin, 377
Canada, 1229
cannabis, 295
cardiac rehabilitation, 889
cardiovascular disease, 52, 150, 208, 311, 810, 1082, 1368
cardiovascular disease risk, 730
cardiovascular risk, 1149
cardiovascular risk factors, 363, 581, 752, 1129, 1274
case-control study, 776

central obesity, 716, 1141 cGMP, 969 Charcot, 1243 Charcot foot, 1334 childhood diabetes, 531 children, 1144 Chinese, 349, 1274 chlorpropamide, 625 cholesterol, 383 chronic complications, 252 chronic painful peripheral neuropathy, 976 chronic wounds, 786 cigarette smoking, 188 cilazapril, 471 cine MRI, 524 circulatory diseases, 936 clamp, 568 classification, 987 clinical score, 1149 clinical trial, 769 cocaine, 295 cognitive function, 230, 440, 999, 1253 cognitive testing, 1366 cohort, 845 cohort study, 685 collagen IV, 1102 colonization, 1247 complex regional pain syndrome, 1334 complications, 91, 203, 657, 1304 compound heterozygosity, 380 continuous glucose monitoring, 487 continuous glucose monitoring system, coronary arterial disease, 342 coronary artery bypass, 790 coronary artery disease, 776 coronary calcium score, 1129 coronary heart disease, 238, 738, 889, 901, cost-effectiveness, 460 costs, 1229 counter-regulation, 440 coverage, 271 Cox regression model, 311 coxsackievirus infections, 507 C-peptide, 349, 354, 428, 1007, 1310 C-reactive protein, 39, 203, 931 cross-sectional study, 363, 798 Crow-Fukase syndrome, 292 CRP, 810 CRPS-1, 1334 cystic fibrosis, 691 cystic fibrosis-related diabetes, 691

DARTS, 1353

database, 1353

death certification, 936

deep tissue biopsy, 705

DCCT, 673

driving, 1014 drug therapy, 666 duration of diabetes, 311 EC 'In Vitro Diagnostic' (IVD) directive, ECG stress test, 342 echocardiography, 524 ecstasy, 295 education, 122, 599 elderly, 677 elderly patients, 999 electrical stimulation, 940 endothelial dysfunction, 171, 262 endothelial function, 931, 983, 1082 endothelium, 552 enterovirus, 507 enteroviruses, 156 epidemiology, 103, 716, 906, 1041, 1353 epilepsy, 628



equations, 318 erythromelalgia, 1334 erythropoietin, 1174 Escherichia coli, 1032 ethnicity, 388, 716, 1090 evaluation studies, 1066 exercise, 122

faecal incontinence, 176 familial clustering, 358 familial hyperinsulinaemia, 628 family practice, 1221 fasting plasma gulcose, 874 fat localization, 881 fieldwork, 1353 first-degree relatives, 262 fluid retention, 945 foot care, 165 foot ulcer, 1334 foot ulceration, 1161 foot ulcers, 987 Framingham, 318 free fatty acids, 171 functional status, 793

gangrene, 987 gastric emptying, 1204 gastrointestinal absorption, 354 gene, 1325 gene polymorphism, 1113 genetics, 752, 837 geographical distribution, 256 gestational diabetes, 68, 388, 829 gestational diabetes mellitus, 103, 1181 gestational, 1310 GIS, 256 glargine, 1213 glibenclamide, 859 glitazones, 666 glucagon, 1204 glucose, 39, 782, 1102, 1280, 1346 glucose clamp, 1090 glucose-clamp, 763 glucose control, 1253 glucose profiles, 468 glucose-lowering treatment, 896 glutamic acid decarboxylase autoantibodies, 559 glycaemic control, 311, 460, 491, 657, 896, glycated haemoglobin, 657, 1200 glycosylated haemoglobin, 691

haematocrit, 476 haemodynamic parameter, 223

Graves' disease, 267

guideline adherence, 586

Greenland, 1237

group care, 644

gut flora, 64

haemoglobin A₁₀, 129 haemoglobin variants, 1041 haemoglobinopathies, 377 haemoglobinopathy, 1041 HbA₁₀, 305, 377, 468, 657, 673, 677, 874, 1041, 1263, 1304 HDL, 423 health education, 1274 health insurance, 45 health outcomes, 592 health status, 45 health-related quality of life, 1075 heart, 183 heart disease, 318 heart failure, 945 heart rate, 1025 heathcare costs, 1269 hemiplegia, 623 heparan sulphate, 1135 hepatocyte nuclear factor-1 alpha, 928 heroin, 295 high-sensitivity C-reactive protein, 336 HMG-CoA reductase inhibitors, 962 Ho Chi Minh City, 371 home management, 531 homeostasis model assessment, 136 homeostatic model assessment, 568 honeymoon period, 1316 hospital demand, 32 hospital inpatients, 1269 HPLC, 377, 1041 hygiene hypothesis, 1035 hypercholesterolemia, 599 hyperglycaemia, 176, 246, 577, 859, 969 hyperinsulinemia, 746 hyperlipidaemia, 150 hypertension, 150, 165, 383, 471, 599, 823, 1181 hypertriglyceridaemia, 380 hypoglycaemia, 230, 440, 487, 623, 769, 1014, 1366 hypoglycaemia awareness, 487

IDDM2, 267 IFCC HbA_{1c} reference method, 673 image analysis, 84 imaging, 1280 impaired fasting glucose, 371, 1346 impaired fasting glycaemia, 363 impaired glucose homeostasis, 730 impaired glucose tolerance, 363, 403, 415, 1229 incidence, 311, 476 inequality, 45 infection, 705, 987 inflammation, 39, 203, 581, 810, 962, 1249 inhaled insulin, 763 inheritance, 759 insulin, 623, 962, 1014, 1263, 1310

insulin action, 1090 insulin analogues, 1213 insulin antibody, 1316 insulin aspart, 171, 769 insulin autoantibody, 1316 insulin deficient, 1007 insulin lispro, 279 insulin resistance, 52, 103, 188, 203, 208, 262, 515, 568, 628, 859, 906, 962, 1141, 1149, 1237 insulin resistance syndrome, 39 insulin secretagogue, 329 insulin secretion, 73, 1090 insulin sensitivity, 59, 415 insulin therapy, 999 insulin treatment, 1298 Insulin-dependent diabetes mellitus, 1035 insulin-treated diabetes mellitus, 845 intensive insulin therapy, 279 interference, 782 intravenous drug abuse, 1269 Inuit, 1237 iontophoresis, 428 iron, 798 islet antibodies, 447 isoprostanes, 285

Japanese men, 476

ketone sensor, 782 kidney function, 1174 Korea, 1141

laser Doppler velocimetry, 428
laser treatment, 26
latent autoimmune diabetes in adults, 456
LDL oxidation, 142
left ventricular diastolic dysfunction, 524
left ventricular hypertrophy, 823
linkage and association, 218
lipid-lipoprotein profile, 730
lipid-lowering, 901
lipid-lowering drugs, 666
lipid-lowering drugs therapy, 1108
lipids, 329
lipoprotein lipase, 380, 1135
lipoproteins, 142, 423
longitudinal study, 311
long-term Type 1 diabetes, 539

macrophage migration inhibitory factor, 1292 malignant melanoma, 924 management, 165 mass screening, 434, 1066 matrix metalloproteinases, 1102 maturity-onset diabetes of the young, 928 medical management, 666 MEMO, 1353 memory, 1253



mental retardation, 628 meta-analysis, 18, 114, 609 metabolic control, 1253 metabolic remission, 1316 metabolic syndrome, 52, 252, 363, 906, 1141, 1237, 1339 metformin, 136, 829, 1200 microalbuminuria, 26, 246, 383, 1192 microangiopathy, 837 microbial flora, 1035 microbiological analysis, 705 microflora-associated characteristics, 64 micronutrients, 677 microvascular complications, 867, 1325 mixing insulin, 604 MODY, 928 monocyte chemotactic protein-1, 1292 monounsaturated fatty acids, 142 mortality, 183, 305, 790, 845, 1243 mortality rates, 936 multidrug-resistant microorganism, 710 multiple sclerosis, 267 mutation screening, 218 myocardial infarction, 183 myocardial metabolism, 1280 myocardial oxygen demand, 1025

national survey, 677 nephropathy, 852 neural network, 84 neuropathic foot ulcers, 786 neuropathy, 91, 183, 852, 987, 1243, 1334 Neuropathy Impairment Score, 114 neuropsychological sequence of hypoglycaemia, 1366 neurothesiometer, 563 newly diagnosed diabetes, 896 nifedipine, 1192 nitric oxide, 68, 262, 969 nitrosative stress, 577 nitrotyrosine, 577 nocturnal hypoglycaemia, 1213 non-hypertensive, 1192 nutrition, 122

nateglinide, 983

obesity, 39, 77, 122, 383, 752
offloading, 1161
older adults, 1185
olive oil, 142
ophthalmoscopy, 434
optical density, 604
optometrist, 632
oral antidiabetic drug, 329
oral glucose tolerance test, 581, 691
oral hypoglycaemic, 1082
oral hypoglycaemic drugs, 685
oral insulin, 354
osteoporosis, 1020

osteoprotegerin, 1334 oxidative stress, 423, 552, 577, 1249,

PAH, 928 painful diabetic neuropathy, 940 Pakistan, 920 paraoxonase 1, 837 parents' experience, 531 patient compliance, 129 patient education, 460, 644 peak VO2, 889 pen injectors, 604 perceived health status, 992 percutaneous coronary, 790 perindopril, 1192 peripheral arterial disease, 724 peripheral vascular disease, 91, 336, 987 peroxidation, 68 peroxynitrite, 68 pH, 782 phage display, 1316 pharmacokinetic, 763 phenotypic markers, 928 phenylalanine, 928 physical activity, 77, 615 physical training, 59 physician practice patterns, 150 pilot studies, 1066 pioglitazone, 136, 568, 859 plasma glucose, 305 platelet, 68 plethysmography, 1298 POEMS syndrome, 292 point-of-care systems, 1288 Poisson, 914 polycystic ovary syndrome, 829 polymorphism, 837, 1325 polymorphism (genetics), 1310 Polynesians, 122 population-based study, 752 positron emission tomography, 1280 post-challenge glucose, 581, 636 post-challenge glucose metabolism, 983 postprandial, 279 postprandial glucose, 208 postprandial glycaemia, 468 post-prandial, 1200 post-prandial hyperglycaemia, 171 postprandial lipids, 208 PPARy, 1249 practice guidelines, 150 pramlintide, 1204

pre-eclampsia, 829, 1181

pre-prandial, 1200

preprandial glycaemia, 468

prevalence, 45, 371, 640, 738, 976

primary open-angle glaucoma, 609

primary prevention, 122, 238, 403

primary care, 32, 592, 640, 685, 1096

private sector, 45
pro-inflammatory cytokines, 552
proliferative diabetic retinopathy, 559
proliferative retinopathy, 852
prospective design, 324
prospective study, 1185
protein kinase C β, 1113
protein kinase C-β inhibitors, 666
psychological well-being, 992
psychosocial factors, 1263
puberty, 246
pulsatility index, 223

quality, 32 quality assurance, 1066, 1121, 1288 quality of life, 644

randomized clinical trials, 644 randomized controlled trial, 491, 592, 1082 randomized trial, 901 RANK-L, 1334 reactive hyperaemia, 931 readiness to change, 122 reflex sympathetic dystrophy, 1334 regression models, 914 regular insulin, 171 remission, 447 renal function, 1096 repaglinide, 1200 resistance, 1032 resistive index, 223 retinopathy, 165, 632, 852 rhinoviruses, 156 risk, 318 risk assessment, 238 risk factor, 423, 724 risk factors, 103, 371, 1096 risk score, 874 rodent, 759 rosiglitazone, 415, 1280 RSD, 1334 Rydel-Seiffer tuning fork, 563

sagittal diameter, 1339
schizophrenia, 515
screening, 84, 271, 456, 632, 874, 992, 1075, 1121
second derivative of photoplethysmogram, 636
secondary care, 640
secondary diabetes mellitus, 1049
self-management control, 491
self-monitoring, 1045
sensitivity, 874
severe hypoglycaemic, 914
short-chain fatty acids, 64
sialic acid, 203
sickle cell trait, 483



silent myocardial ischaemia, 342 single slice MRI-scan, 1339 slit-lamp biomicroscopy, 632 SNPs, 1113 social service, 999 socioeconomic factors, 129 sodium reabsorption, 969 soleus muscle, 539 somatostatin analogues, 666 South Asian, 388 South Asian ethnicity, 1357 South Asians, 845 specialist general practitioners, 32 specificity, 874 spironolactone, 471 staged diabetes management, 592 statins, 1020 stiffness, 539 stool samples, 156 street drugs, 295 stress hyperglycaemia, 305 stretch reflex, 539 stroke, 623 Strong Heart Study, 311 structured care, 1221, 1274, 1357 structured treatment and teaching programme, 999 sudden hearing loss, 96 survey, 271 survival, 936 symptoms of hypoglycaemia, 1014 symptoms of neuropathy, 852 systematic review, 507 systolic function, 524

T helper cell immune reactions 1 and 2, 1035

Taiwanese, 724 TDT, 218 telemedicine, 1121 thallium-201 scintigraphy, 342 therapy, 1161 thiazolidinedione, 329 thiazolidinediones, 810, 945, 1249 thiobarbiturate acid reacting substance, 1346 time-action, 763 Tirana, 77 tissue factor pathway inhibitor, 1135 tissue inhibitor of matrix metalloproteinases, 1102 tobacco, 285 Total Symptom Score, 114 trace elements, 256 transferrin saturation, 798 transforming growth factor β₁, 818 transforming growth factor beta 1, 285 transluminal, 790 treatment, 976 treatment of hypoglycaemia, 1014 triglycerides, 171, 208, 615 tropicamide, 434 Type 1 diabetes, 26, 64, 156, 218, 256, 267, 279, 324, 358, 460, 468, 487, 769, 867, 914, 1204, 1269, 1316, 1366 Type 1 diabetes mellitus, 96, 428, 818, 931, 1213, 1263 Type 1 DM, 447 Type 2 diabetes, 52, 59, 73, 77, 103, 122, 188, 208, 252, 262, 311, 336, 568, 644, 685, 724, 752, 776, 810, 837, 859,

920, 983, 992, 1045, 1075, 1090, 1113,

1129, 1174, 1185, 1192, 1253, 1274,

1298, 1325, 1357, 1368

Type 2 diabetes mellitus, 403, 456, 476, 483, 592, 738, 746, 1020, 1144, 1304

UKPDS, 673, 1368 ulcer swabbing, 705 ultrasonography, 223 uptake, 1108 urban-rural difference, 256 uric acid, 724 urine glucose self-monitoring, 685 urine testing, 1045 uropathogen, 1032

vascular adhesion molecules, 262
vascular calcification, 1334
vascular endothelial growth factor, 292,
545, 697
vasodilatation, 428
vasomotor neuropathy, 1334
VEGF, 292
vibration perception threshold, 563
visceral fat, 1339
visceral obesity, 730
vitamin C, 552
vitronectin, 1135

waist circumference, 383, 1339 Warren repository, 358 WHO criteria, 1141 Wolfram syndrome, 218 women, 615, 730

young adults, 447 younger-onset Type 1 diabetes, 559

zero-inflated negative binomial, 914

